

Ann Arbor

STATE OF MICHIGAN

DEPARTMENT OF CONSERVATION

P. J. Hoffmaster, Director

GEOLOGICAL SURVEY DIVISION

R. A. Smith, State Geologist

Publication 40
Geological Series 34
Part I

**The
Pleistocene History of the Tahquamenon
and Manistique Drainage Region of the
Northern Peninsula of Michigan**

STANARD G. BERGQUIST, PhD.
Michigan State College

PREPARED UNDER THE DIRECTION OF
R. A. SMITH, State Geologist



PUBLISHED AS A PART OF THE ANNUAL REPORT OF THE GEOLOGICAL SURVEY
DIVISION FOR 1936.

STATE OF MICHIGAN
DEPARTMENT OF CONSERVATION
P. J. Hoffmaster, Director
GEOLOGICAL SURVEY DIVISION
R. A. Smith, State Geologist

Publication 40
Geological Series 34

OCCASIONAL PAPERS
ON THE
GEOLOGY OF MICHIGAN
PREPARED UNDER THE DIRECTION OF
R. A. SMITH, State Geologist



PUBLISHED AS A PART OF THE ANNUAL REPORT OF THE GEOLOGICAL SURVEY
DIVISION FOR 1936.

FRANKLIN DEKLEINE COMPANY, PRINTERS, LITHOGRAPHERS, BOOKBINDERS, LANSING

*
41213.



LETTER OF TRANSMITTAL

*To the Honorable, the Director and the Board of Commissioners of the
Department of Conservation of the State of Michigan:*

Hon. P. J. Hoffmaster, Director
Hon. William H. Loutit, Chairman
Hon. Philip Schumacher
Hon. Harry H. Whiteley
Hon. M. J. Fox
Hon. Philip K. Fletcher
Hon. Joseph P. Rahilly
Hon. William J. Pearson
Hon. Wayland Osgood, Secretary

Gentlemen: In compliance with Act No. 65 of 1869 as amended by Act No. 179 of 1871 of the Public Acts of Michigan under which the Geological Survey Division of the Department of Conservation operates, I have the honor to present herewith three reports on the geology of Michigan, by Prof. R. C. Hussey of the University of Michigan, Prof. W. A. Kelly and Prof. S. G. Bergquist of the Michigan State College, and recommend that they be published as Publication 40, Geological Series 34, of the Geological Survey Division, Department of Conservation, as part of the Annual Report for 1936.

Prof. Bergquist's report on the surface geology and development of the Tahquamenon-Manistique drainage system was prepared as a part of the Land Economic Survey of the eastern part of the Northern Peninsula and is a study and description of the surface conditions in the drainage basins of the two main river systems of Luce, Alger, and Schoolcraft counties, and the origin of scenic features in the State Forests of these counties. A foreword has been added, briefly outlining the glacial history of the State and the development of the Great Lakes and containing definitions of glacial features. The report with its accompanying maps and illustrations should be of interest and value not only to the student of glacial geology, soil, and forest conditions, but to tourists as well.

The report on the Trenton and Black River rocks of Michigan by Dr. R. C. Hussey was prepared as a part of the routine study of the rock exposures of the State under the plan of Geological Survey activities. The report is of very timely interest at present, not only because of its scientific value but also because of its economic application in the development of the petroleum interests in the State. The Trenton and Black River rocks are almost the lowest rocks in the State in which one may look for oil accumulations. These rocks rise to the surface in the Northern Peninsula and slope southward under the basin in the central part

of the State. Little descriptive matter has been published about them; thus the report of Dr. Hussey is a valuable contribution to the literature of the formation which may in the future solve the petroleum problems of the State. At present the Trenton has been explored only around the edges of the State, but as upper oil "sands" are depleted deeper drilling will become the practice, in search for new supplies. A report such as this will be of value to the companies engaged in the deeper explorations.

The report on the Pennsylvanian system of Michigan was prepared by Dr. W. A. Kelly of Michigan State College (like the report on the Trenton by Dr. Hussey), as a part of the plan of the Survey to continue a comprehensive study of each of the geological formations of the State. The Pennsylvanian is the youngest system of rocks of the State and contains the coal beds, some gas "sands", water zones, and, where the rock comes to the surface, some building stone is obtained. In addition the Pennsylvanian assumes an economic value in the petroleum development of the State. A common practice in oil exploration is the drilling of shallow test wells to some "marker" bed. Because the Pennsylvanian rocks are the upper rocks of the central part of the State and are relatively shallow, they assume great importance in shallow test drilling and a comprehensive description of them is of value to the oil industry as well as to general geology.

The three reports are outstanding contributions of both scientific and economic value and should aid in the development and utilization of the eastern part of the Northern Peninsula as well as aid in the development of the mineral industry of the State and be a marked contribution to the geologic literature of Michigan.

Respectfully submitted,

R. A. SMITH,

State Geologist.

September, 1936



Digitized by the Internet Archive
in 2026

https://archive.org/details/IA41551510_0127

**PART I. THE PLEISTOCENE HISTORY OF THE
TAHQUAMENON AND MANISTIQUE DRAIN-
AGE REGION OF THE NORTHERN PEN-
INSULA OF MICHIGAN**

**STANARD G. BERGQUIST, PhD.
Michigan State College**

THE PLEISTOCENE HISTORY
OF THE TAHQUAMENON
AND MANISTIQUE DRAINAGE
REGION OF THE NORTHERN
PENINSULA OF MICHIGAN.

By
STANARD G. BERGQUIST

A dissertation submitted in partial
fulfillment of the requirements for the
degree of Doctor of Philosophy in the
University of Michigan.

JUNE, 1933

Published as a part of the Annual Report of the Michigan Geological
Survey Division, Department of Conservation, for 1936



CONTENTS

	Page
Foreword, Discussion of Glaciation in Michigan.....	17
Synopsis	41
Introduction	47
Acknowledgments	47
Chapter I—The Tahquamenon Drainage System	49
General Considerations	49
The Two Hearted River System.....	54
The Little Two Hearted River.....	55
The Sucker Basin	55
The Shelldrake Drainage System	55
Lakes	56
Swamps and Marshy Tracts	57
Chapter II—The Manistique Drainage System	59
The Manistique River	59
Chapter III—Glacial Activity	63
Features Developed by the Green Bay Lobe.....	63
The Trenary Till Plain	63
Position within the axis of the Green Bay Lobe.....	63
Topography and Relief	63
Character of the Drift	65
The Eskers in the Trenary Till Plain	66
Drumlins	68
Conclusions	68
The Cook's Moraine	68
Outwash Associated with the Cook's Moraine.....	69
Features Developed by the Superior Lobe	69
Moraines	69
The Newberry or "Outer" Morainic System.....	69
The Steuben Segment	71
The Hiawatha Segment	71
The Blaney Segment	73
The Munising or "Inner" Morainic System	75
Significance	80
The Crisp Point Moraine	81
Outwash Plains	82
Outwash Plains Associated with the Munising Moraine.....	83
The Wetmore Plain	83
The Boot Lake Plain	85
The Kingston Plain	86
Chapter IV—Lacustrine Activity	88
General Discussion	88
Lake Algonquin	89
Shore Features Pertaining to Lake Algonquin	95
Highest Algonquin Shore in the Tahquamenon Region.....	95
Lower Algonquin Shores in the Tahquamenon Region.....	96
The Lowest Algonquin Shore	99
Old Shores in the Grand Marais Region	99
Algonquin Shores in the Manistique Basin	101
The Nipissing Great Lakes	104
The Nipissing Shore Features	104
Lake Superior Region	104
Lake Michigan Region	106
Post-Nipissing Shore Features	108
The Algoma Shore Features	110

	Page
Chapter V—Aeolian Activity	113
Shore Dunes Along Lake Superior.....	113
Alger County (Shore Dunes)	113
The Grand Sable Bluffs	119
Luce County (Shore Dunes)	121
Dunes of the Poorly Drained, Marshy Lowland Plains of the Main Drain- age Ways	122
The Seney Complex	125
The Manistique Complex	125
Inland dunes	129
Shore Dunes Along Lake Michigan.....	130
Summary and Conclusions	132
Glacial activity	132
Lacustrine activity	133
Aeolian activity	135
Bibliograhny	139

ILLUSTRATIONS

FIGURES

	Page
Figure 1. Map of North America showing glaciated area	18
Figure 2. Map of Glacial lobes	22
Figure 3. Map showing distribution of moraines	24
Figure 4. Map showing development of Lake Maumee	29
Figure 5. Map showing development of Lake Whittlesey	31
Figure 6. Map showing development of Lake Warren	32
Figure 7. Map showing development of Lake Lundy	33
Figure 8. Map showing development of Lake Algonquin	34
Figure 9. Map showing development of Lake Nipissing	37
Figure 10. Map showing hinge lines and present Great Lakes	38
Figure 11. Map showing Physiographic Provinces of the Northern Peninsula with Location of the area included in the Report	42
Figure 12. Geological Map Showing Surface Formations of Luce County.....	53
Figure 13. Geological Map Showing Surface Formations of Schoolcraft County	58
Figure 14. Geological Map Showing Surface Formations of Alger County.....	64
Figure 15. Geological Map Showing Surface Formations of the Tahquamenon and Manistique Drainage Basins, including Alger, Luce, and Schoolcraft Counties	137
Figure 16. Location Map Showing Major Features of the Tahquamenon and Manistique Drainage Basins, including Alger, Luce, and Schoolcraft Counties.....	137

LIST OF PLATES

Plate	Fig.	Page
I.	1. The Tahquamenon River above the Upper Falls.....	50
	2. The Upper Falls of the Tahquamenon River.	
II.	1. Sandstone Gorge below the Upper Falls of the Tahquamenon River.....	52
	2. Looking north across the Tahquamenon Basin from the crest of the Niagaran Escarpment, just north of Hendricks Quarry.	
III.	1. East Branch of the Fox River, one of the headwaters of the Manistique River, in a muskeg swamp	60
	2. The Fox River, near its headwaters, valley incised in a sandy outwash plain.	
IV.	1. Portion of Trenary till plain north of Trenary. Esker ridge in background	64
	2. Esker ridge on Trenary till plain, near Limestone.	
V.	1. Drumlin in Till Plain, ½ mile west of Eben, Alger County.....	67
	2. Cross section of drumlin showing heterogeneous arrangement of boulders and cobble in till.	
VI.	1. A portion of the Steuben section of the Newberry Morainic System. Fire Tower hill, near Steuben	70
	2. A closer view showing undulating character of topography.	
VII.	1. Morainic topography in the vicinity of Blaney. A portion of the Newberry Morainic System shows characteristic level relief of water worked feature	72
	2. Water worked morainic area of the Newberry System in the Twin Lakes country.	
VIII.	1. High morainic bluff in the Munising System, near the city of Munising..	76
	2. Moraines of the Newberry System, near Laketon.	
IX.	1. Burned over outwash plain, South of Kingston Lake.....	84
	2. Section of outwash plain showing stratification in the sandy material.	
X.	1. Algonquin terrace and wave-cut cliff, 1 mile south of Chatham.....	90
	2. Upper Falls on the Au Train River, near Forest Lake.	
XI.	1. Algonquin shore ridge, at 870 feet A. T., against the remnant of morainic ridge, Luce County	94
	2. Closer view of shore shown in Fig. 1.	
XII.	1. Algonquin terrace and shore cliff. Nipissing scarp in extreme left. Near Grand Marais	98
	2. Nipissing shore scarp and terrace. Near Grand Marais.	
XIII.	1. Algonquin shore cliff. Near headwaters of Fox River.....	100
	2. Algonquin shore scarp at entry with Fox River Valley. 4½ miles north of Seney.	
XIV.	1. A view into Sturgeon Hole, a portion of the Manistique swamp, through the rock corridor in the S. ½, Sec. 14.....	102
	2. Limestone block pavement on crest of a water washed moraine. Most of the till has been removed by water activity.	
XV.	1. Section through a cobbly, gravelly shore showing stratified layers.....	103
	2. A closer view of the feature showing the coarse assortment of material, mostly limestone.	
XVI.	1. Nipissing terrace with shore cliff in background. Two miles east of Manistique	105
	2. Sand dunes on Nipissing plain. Ponded slough of Manistique River in background.	
XVII.	1. Wave-cut Nipissing rock cliff. City of Manistique	107
	2. Nipissing terrace and wave-cut shore cliff. ¾ mile south of Thompson.	
XVIII.	1. Sandy lake plain on crest of the Nipissing escarpment, 2 miles east of Manistique	109
	2. Algoma shore ridge exposing rubble structure. Manistique.	
XIX.	1. Gently dipping limestone floor along shore of Lake Michigan. South of Seul Choix Point	111
	2. Glacial erratics on limestone lake floor. Along shore of Lake Michigan just east of Manistique.	

Plate	Fig.	Page
XX.	1. Crest of Grand Sable dunes showing deflation hollows. Near Grand Marais	114
	2. Exhumed forest on crest of wind-swept sand dunes. Near Grand Marais. Lake Superior in background.	
XXI.	1. Perched dunes resting on cobbly bar 275 feet above level of Lake Superior. Grand Sable area	116
	2. Sand dunes encroaching upon channel of Grand Sable Creek. Near Grand Marais.	
XXII.	1. A lag-gravel plain with perched sand dunes in background. Grand Sable dune area, near Grand Marais	117
	2. Mud streams on north flank of Grand Sable dunes, near Grand Marais.	
XXIII.	1. Sand bluffs near Au Sable Point	120
	2. Sand dunes in the Sucker River area.	
XXIV.	Aerial photograph of Schooner county showing dune ridges.....	124
XXV.	1. A portion of the Manistique swamp. Near Walsh.....	127
	2. Sand dune topography near Seney as viewed from the Fire Tower.	
XXVI.	1. Dune blown up against the slope of a moraine in the Manistee swamp...	128
	2. Section of dune showing cross-bedding in the unconsolidated sand. Tahquamenon swamp, 2 miles north of McMillan.	
XXVII.	Aerial photograph of lower Manistique Basin.....	136

FOREWORD

THE GLACIAL HISTORY OF MICHIGAN

Centers of Ice Accumulation

During hundreds of millions of years great seas invaded much of the continent of North America and vast accumulations of gravel, sand, mud, and lime were deposited in these basins to form the foundational rocks out of which Michigan was developed. But the actual sculpturing of the present surface, the development of the hills and valleys, or relief, and the delineation of the State into the definite physical unit we know today was not accomplished until yesterday in geological time, rather late in the Pleistocene or glacial epoch.

During this rather recent event, vast sheets of ice in the form of glaciers gathered in the snowfields of Canada and slowly spread over the surface of the northern half of the North American continent, covering its surface with a blanket of ice that was probably a mile or more in thickness. There were four major and several minor centers of refrigeration and ice accumulation in North America during this time. (Fig. 1). One center which was on the plateau east of James Bay is named the Labradorean. The greatest ice movement was southwestward from this plateau. The ice advanced 1,600 miles almost to the present Ohio River, whose course was more or less definitely established along the border of the glacier. Locally the ice advanced slightly into southern Illinois. It was this mass of ice that spread over Michigan in several invasions and shaped the hills, valleys, and plains and placed the rivers and lakes almost as we know them today. And it was the melting water from this ice which collected in basins to form the Great Lakes.

Another glacier, the Keewatin, developed on the low, flat lying plains west of Hudson Bay. It was the largest of the three ice fields and in its greatest movement spread southward nearly 1,500 miles, to the position of the Missouri River, whose course was unquestionably defined along its margin.

The third center, the Cordilleran, occupied the basin-like plateau between the Selkirk Mountains and the Cascade Range. Its activity was confined largely to the lofty mountains of western North America with little or no movement into the surrounding lower regions.

Greenland was also a center of accumulation. Here the ice spilled over the borders of the continent into the surrounding seas but did not join the main ice mass of the North American Mainland.

During the greatest ice activity in the Pleistocene epoch the continental glaciers covered approximately 4,000,000 square miles of surface in North America, 2,000,000 square miles in Europe, another 2,000,000 square miles in Asia, 5,000,000 square miles in Antarctica, and 800,000 square miles in Greenland.

Today, the ice sheets have largely melted from North America, Europe and Asia, but still remain in Antarctica and Greenland covering practically all the south polar area and all but the marginal tracts of Greenland.



Figure 1.—Map of North America showing the area covered by ice during the Ice Age. After W. C. Alden.

Ice Invasions

In spite of varied controversies regarding certain details of activity, glacialists are generally agreed that the ice blankets melted away and

then reaccumulated several times causing renewed invasions of the ice sheets during the Pleistocene epoch. These invasions were separated from each other by long intervals of time during which the glaciers melted back into the accumulation centers.

For convenient reference each ice invasion has been given a specific name and each interval of time between invasions—interglacial stage—has also been named, referring to those places where the glacier left important records which glacial geologists have learned to read.

These records consist of:

1. Long lines and festoons of hills—The Moraines. The rock debris which the ice had scooped off the land in its steady push out from the various centers and which was left behind when the ice melted comprise the *moraines*. If the rate of backward melting was equal to that of the forward push, the front of the ice remained stationary for relatively long intervals and the load of glacial drift was dumped along its front in ridges of rounded hills interspersed with valleys and basins—the *marginal moraine*. Each moraine in the state has been given a name, usually the name of a town or river which is an important geographic location on the moraine.

2. Ground Moraines—formed when rapid recession of the ice, due to excessive melting, caused the rock waste to be spread out over the surface from which the ice had just retreated. The moraines are constructed of all types of debris that the ice was able to collect, including boulders, cobbles, gravel, sand and clay, intermixed and without assortment. The ground moraines, on the other hand, are constructed largely of un-assorted clay and sand with scattered boulders. They are usually fairly level and smooth in relief and are known also as *till plains*.

3. Outwash or Overwash Plains—formed by rivers flowing out from the edge of the melting glaciers washing the gravel, sand and clay from the ice and spreading them out in front of the moraines. In the sorting process the coarse material such as cobble and gravel was deposited close to the ice margin, and the finer material such as sand and clay was carried farther out for deposition. Each time the progress of ice retreat was halted, moraines and outwash plains were built up. Whenever the ice melted back rapidly with no long stops in recession, *ground moraines* were developed. When river channels in the ice became choked with sand and gravel, *eskers* were left.

4. Eskers—long narrow ridges commonly referred to as hogs backs and Indian trails. Rivers existed in the ancient glaciers as they do in the glaciers of Switzerland and Alaska today. The river channels became filled with the sand and gravel in the ice, which, when it melted, allowed the channel shaped masses of assorted sediment to settle gently to the underlying surface of bed rock or till plain.

All the glacial deposits taken together are known as *drift*, and by means of the drift the great glacier wrote its own story for future students to read just as the glaciers of today are leaving similar but less extensive records for modern study.

Nebraskan Invasion

The earliest of the glacier invasions is commonly referred to as the Nebraskan; it started approximately a million years ago or about the time that the Ape-man of Java (*Pithecanthropus erectus*), came upon the earth. This glaciation was undoubtedly the most extensive as well as the longest of the several stages.

The Nebraskan invasion was followed by the Aftonian interglacial stage during which the glaciers melted back to near the centers of accumulation.

Kansan Invasion

The second invasion of the ice sheets was the Kansan which, according to Leverett, dates back some 450,000 to 600,000 years ago. During this stage the ice reached southward as far as the northeastern part of Kansas and covered all but the outer edges of the Nebraskan glacial deposits, the so-called gumbo-tills in the region to the west of the Mississippi River. The ice of this invasion advanced from the Keewatin center and passed over Manitoba, Minnesota, Iowa and Missouri and encroached broadly on the Dakotas, Nebraska and Kansas.

Then followed another interval of melting, the Yarmouth interglacial, and the Kansan drift surface was exposed to pronounced weathering. Deposits of peat, and forest beds 15 feet thick lying above the Kansan drift give evidence of a relatively long interval of ice removal. It is probable that the pre-historic Heidelberg man appeared in Europe during the time of the Yarmouth interglacial stage.

Illinoian Invasion

The third invasion, the Illinoian, occurred between 150,000 to 200,000 years ago, and was marked by a decided readvance of the ice from the Labradorean center. This time, the ice spread across the district of the Great Lakes, westward as far as the Mississippi and southward approximately to the line of the Ohio River. The Illinoian drift is exposed in southern Ohio and northern Kentucky where it shows the effects of leaching by the action of weather and water to a depth of two to five feet. In Michigan and Wisconsin the Illinoian drift is everywhere almost completely buried under drift of the last or Wisconsin invasion. It is exposed in the bed of the Raisin River at Monroe and in numerous stream gulleys along the shore of Lake Huron in the Thumb region. Wells

drilled through the glacial drift usually encounter the hard, indurated, blue clay of the Illinoian directly beneath the brownish surface, or Wisconsin, till and immediately above bed rock in many portions of the state. Because of its hardness due to compaction by ice pressure, it is frequently mistaken for bedrock.

The Sangamon interglacial stage followed the Illinoian invasion. The Sangamon soil, composed of a dark colored gumbotil contains much organic matter and is usually buried under deposits of a very fine wind deposited silt soil commonly known as *loess*. The Sangamon gumbotil, ranging in thickness from three to five feet where best preserved, seems to bear evidence of having been developed under conditions which suggest a cool, damp climate.

Wisconsin Glaciation

The last invasion of the ice sheets from the centers of refrigeration is known as the Wisconsin glaciation. In the early part of the Wisconsin stage the ice radiated from the Labradorean center and spread southward across the Great Lakes region, into central Illinois and Indiana. When the ice wasted away by melting it deposited the brown drift so characteristic of Michigan. In the later Wisconsin stage the ice from the Keewatin center spread over Minnesota, Iowa and the Dakotas and laid down a considerable volume of gray drift which in places overlaps the brown drift of earlier date.

On the basis of the amount of weathering developed in the surface of the latest drift, Leverett estimates that the last remnants of the Wisconsin ice sheet left the southern tier of counties in Michigan not more than 35,000 years ago. By the same measure it is probable that the final vestiges of the ice were removed from the basin of Lake Superior some 10,000 years ago, or about the time that Cromagnon civilization was spreading in Europe.

Throughout the hundred of millions of years which followed the removal of the last, the Paleozoic, seas from Michigan and up to the beginning of the glacial epoch about a million years ago, the exposed bed rock surface of the state was subjected to widespread destruction and up-building by rivers, winds, rains, and frost or a prolonged *erosional* activity. The pre-glacial surface was cut almost to a plain, or a vast peneplain scored by old rivers which had carved their valleys deep into the rock floor. Because of the thick cover of glacial drift which now mantles the bed rock surface it is difficult to accurately reproduce the pre-glacial drainage of the Great Lakes region. Scattered deep well records, however, seem to indicate that before glaciation there were a number of well defined, deeply cut river valleys following in part the lines which now are occupied by the Great Lakes. Speculation is rife

among glacialists and hydrologists as to the direction of drainage of the pre-glacial rivers. One school of investigators seems to favor a southward drainage into the Mississippi river. Another group is equally convinced that the St. Lawrence Valley was then, as now, the major line of discharge into the Atlantic. Still a third group would have the waters draining northward through some of the major waterways into Canada, with an outflow into the Arctic ocean. Much more subsurface research is necessary to conclusively prove this perplexing problem. In spite of the differences of opinion concerning pre-glacial drainage outlets, the general consensus of opinion is that the major lines of river development were concentrated in troughs which were, at least generally, parallel to the present basins of the larger lakes.




Figure 2.—Map of the Glacial Lobes.

In the Wisconsin stage of glaciation, the Labradorean ice sheet re-advanced southward across the region of the present Great Lakes. As it encountered the free drainage ways of the area, the deeper valleys served as natural thoroughfares for ice flow. The ice front then formed several large lobes and moved forward under control of the river courses. The largest of the ice lobes ploughed through the valley in which the

basin of Lake Michigan was eventually sculptured. This glacier is referred to as the Lake Michigan lobe and in its maximum extension it completely filled the basin and overflowed its rim. It expanded also southward beyond the site of Chicago to form the extension referred to as the Illinois lobe.

A second important mass of ice, the Huron-Erie lobe, was directed through a series of valleys which were later sculptured into the basins of lakes Huron and Erie. A third mass of smaller dimension was pushed out from the Huron-Erie ice extension into the basin of Saginaw Bay and became active as the Saginaw lobe. As this ice mass moved southward it nosed gradually out of the Saginaw drainage depression and onto the upland where it spread out as a relatively thin sheet of ice which eventually extended into the valley of the Kankakee River in northern Indiana. (Fig. 2).

Later still, after the glacier had left the surface of the southern Peninsula, a large lobe became centered in the Lake Superior basin. To the activities of this lobe, the northern Peninsula owes most of its glaciated characters, some of which are described in the discussion of the features of the Manistique-Tahquamenon drainage area.

Development of the Moraines

The Saginaw lobe was initiated in the Saginaw valley as a definite tongue extending out from the much larger Huron ice mass. In its greatest expansion this body of ice reached into the northwestern part of Indiana where evidences of its activity may be traced to a point somewhat south of South Bend. The Saginaw lobe moved through and beyond the present Saginaw Bay and, in the climax of its development, was confined between the Michigan and Huron-Erie lobes with which it joined during the height of the Wisconsin stage. (Fig. 2).

A study of the lines of hills, the morainic systems, distributed over the surface of northern Indiana and southern Michigan, seems to show that the ice of the Saginaw lobe melted back from the basin of the Kankakee River, disappeared completely from northern Indiana and withdrew into southern Michigan, while the Michigan and Huron-Erie lobes still filled to overflowing their respective basins.

Until the Saginaw ice began to melt back in the early Wisconsin stage, the three lobes were welded into a more or less undifferentiated mass. The early retreat of the ice in the Saginaw tongue, however, allowed the various lobes to acquire distinctive and individual characters and to build moraines at their borders. It appears also that the Saginaw lobe because of its thinner ice was unable to transport as large a load of drift as the bordering basin filled lobes, and therefore melted faster. Clean ice,

or ice containing a small amount of drift material, melts faster than dirty ice where the included rock material acts like rock wool, that is, as an insulator preventing melting.

The various halts of the Saginaw lobe as it melted across Michigan are evidenced in the succession of more or less parallel moraines which are found in festoons closing in on Saginaw Bay from the southwest. (Fig. 3). In Indiana and southern Michigan, the moraines of this lobe



Figure 3.—Map showing the distribution of moraines in Michigan.
(After Leverett).

are much more strongly developed than they are farther north. It is probable that in the area of convergence of the three ice tongues in northern Indiana and southern Michigan, a considerable amount of drift

was carried in by the Michigan and Huron-Erie lobes and loaded into the Saginaw ice for later deposition upon its retreat. The excessive loading of the southern end of the Saginaw tongue resulted in its becoming more sluggish which consequently allowed for slower retreat and the development of stronger morainic ridges. Farther north, on the other hand, the influence from the adjoining lobes except in the inter-lobate areas was insignificant and the resulting moraines are rather weak and poorly developed.

In its retreat from the Maxinkuckee moraine which marks the southernmost position of the oscillating front of the Saginaw ice in Indiana, the lobe made a succession of short halts before finally coming to rest on the Kalamazoo moraine. These short stops in the back step of the ice front are marked by a series of moraines, named in order the Bremen, New Paris, and Middlebury in northern Indiana, and the Sturgis and Tekonsha in southern Michigan. (Fig. 3.). These moraines were later partially modified by the lateral extension of the ice sheets from the Michigan and Huron-Erie basins and thus became greatly altered in relief. The first portion of Michigan to be uncovered was the St. Joseph River valley in southeastern Cass and southwestern St. Joseph counties when the Saginaw ice front rested on the Sturgis moraine.

When the retreating Saginaw ice lobe finally came to rest on the position of the Kalamazoo moraine, it stopped for a sufficiently long time to build up an extensive ridge. This moraine, extending from Hastings south and east through Kalamazoo, Marshall, and thence to Devil's Lake in Lenawee County, joins on the east with the Mississinawa moraine which outlines the outermost position of the Erie lobe in Michigan. On the west side of the state, a limb of the Kalamazoo moraine extends from Hastings southward through Kalamazoo and Cassopolis and is in a belt more or less parallel to the present shore of Lake Michigan. It marks the eastern boundary of the Lake Michigan ice lobe.

From the position of these three joined moraines, it is obvious that the Saginaw ice front had melted northward to expose a considerable portion of the surface area of St. Joseph, Branch, Kalamazoo and Calhoun counties while the ice of the Michigan and Huron-Erie lobes still covered the borders of the State. When the ice borders stood on the Mississinawa and outer Kalamazoo moraines, the main drainage of the State seems to have followed the course of the Kalamazoo River westward to the site of the present city of Kalamazoo. Thence, it was directed southward into the St. Joseph Valley past Three Rivers and into the Kankakee Valley near the site of South Bend. The water in this valley was dammed back in Indiana by the Marseilles moraine behind which it was ponded into a large lake, the Kankakee torrent, which eventually grew in places to a width of ten and more miles.

Renewed melting in the Saginaw lobe caused the ice front to retreat to the position of the Charlotte moraine. This moraine connects with the Valparaiso morainic system of the Lake Michigan lobe and may be traced from near Grand Rapids eastward through Charlotte, and Brighton to Milford. The outermost ridge of the Charlotte moraine connects with a ridge of the Valparaiso system at Dias Hill, about 12 miles south of Grand Rapids. A second ridge connects with the Valparaiso system at the bend of the Grand River directly north of Grand Rapids. A third ridge connects near Cedar Springs—some fifteen miles north of Grand Rapids. On the east side of the state near Pontiac, the Charlotte moraine connects with the Fort Wayne-Wabash systems which were formed at the same time by the Huron-Erie lobe.

At the time when the Charlotte moraine was forming, the melt water drainage from the ice in the vicinity of Milford followed the early valley of the Huron River northward to Portage Lake, near Pinckney. Then it continued westward to the Grand River valley, just north of Jackson, which it followed to Eaton Rapids. From here the water was carried through a swampy depression to the vicinity of Charlotte where it entered the valley of Battle Creek which was cut transversely across the Kalamazoo moraine.

A block of stagnant ice which covered the area extending from the site of Eaton Rapids to the vicinity of Charlotte blocked the Grand River valley and prevented discharge northward. The drainage entered the Kalamazoo River where Battle Creek now stands whence it was directed westward to the vicinity of Plainwell. From Plainwell it appears to have passed southwestward through the low area between the Kalamazoo moraine and the receding ice border to the vicinity of Dowagiac. Here a narrow lake (Lake Dowagiac) extending to the site of South Bend, Indiana, was formed in the valley of the present Dowagiac Creek. From this lake, the waters drained out through the Kankakee Valley, into the Illinois River and thence to the Mississippi.

After the formation of the Charlotte moraine and its correlatives, the Valparaiso and Fort Wayne Morainic systems, the ice along all fronts began another retreat. The next stop of the Michigan lobe and the last stop made by it south of Manistee before it retreated into the Lake Michigan basin is represented by the Lake Border morainic system, the ridges of which closely follow the margin of the lake. The Saginaw lobe correlatives of this system are the West Branch and Gladwin moraines north of the Grand River valley and a deployed group of slender moraines south of the Valley. Named in succession from the vicinity of Lansing northward, the moraines of the deployed group are: Lansing, Grand Ledge, Ionia, Portland, Lyons, Fowler, St. Johns, Flint, and Owosso. These moraines are arranged in more or less parallel concentric bands

around Saginaw Bay and represent individually the various short halts made by the Saginaw lobe in its retreat from the Charlotte moraine to the position of the Port Huron moraine. The Huron-Erie correlative of the Lake Border moraine includes the Defiance, Birmingham, Mount Clemens, and Emmett ridges on the east side of the state. These ridges, in places, have been so completely modified by wave action of the later glacial great lakes that it is difficult to trace them in their entire extent.

It is significant to note that the development of the Grand River in its course from near the site of Grand Rapids, eastward to its confluence with Maple River, was dependent entirely upon the successive retreatal stops of the Saginaw lobe as it gradually back-stepped into the Saginaw Bay. During the interval when the moraines of the deployed group of moraines was being laid down, the Grand River channel was a shallow and poorly defined drainage crease in the surface of the glacial drift. Melt waters from the retreating glaciers flowing as border drainage along the outer margin of the ice sheet, especially during the intervals when the slender moraines north of the valley were being constructed, contributed appreciably to the volume of flow in the newly forming Grand River. Thus the channel was extended and deepened progressively as the Saginaw ice melted. The development of the Looking Glass River as border drainage along the margin of the ice when the Portland moraine was forming increased the volume in the Grand River and was to a large degree responsible for the early development of its present valley.

A further recession and subsequent long halt of the Lake Michigan and Saginaw-Huron lobes was responsible for the formation of the Port Huron morainic system extending from Port Huron and Saginaw Bay, north and westward to Grand Traverse County. This feature which forms the backbone of the glacial structures in the state was constructed when only the borders of Michigan were covered by ice. This strongly developed morainic system shows evidences of a long halt of the ice sheet before it finally left the surface of the Southern Peninsula in the late Wisconsin stage.

The last stand of the ice, as it was retiring from the surface of the Southern Peninsula and before it retreated into the Straits of Mackinac, is represented by the slender Cheboygan moraine which is well marked along the shore of Lake Huron immediately west of Cheboygan.

THE DEVELOPMENT OF THE GREAT LAKES

No story of the glacial history of the state would be complete without giving some consideration to the development of the Great Lakes. Michigan is situated in the center of the area of ice activities which caused

the initiation, successive changes, and final growth of the lakes. As has already been stated, the Great Lakes basins occupy the sites of pre-glacial rivers. Into these old and established drainage depressions the continental ice mass sent out its various tongues or lobes which scoured and deepened the valleys into broad troughs.

The Great Lakes were first formed as small lakes in front of the glacier. As the tongues of ice in these broad troughs melted, the melt water collected in those parts of the troughs which were in front of the ice, thus forming lakes which were hemmed in behind morainic ridges, with shores in ice on the glacierward side. As long as the waters did not rise above the moraines or find an outlet through them, the lakes persisted. Like lakes of the present time these old glacial lakes produced shore lines, beaches, sand bars and other lacustrine features which can be traced on the present land surface. As lakes of today dry up and leave their shore lines and beaches above former water levels, so the water of the glacial lakes was drained out and left traces of former shores and beaches for us to read today. The length of time the glacial lakes existed is shown by the size and "strength" of old lake shores and beaches. Because of their level relief and relative width, the glacial lake beach ridges in Ohio, Indiana, Illinois, Wisconsin and Michigan are often utilized as sites for highway construction. As the oscillating ice front melted farther and farther back toward the centers of radiation more of the troughs were exposed and filled with melt water and the lakes increased in size. They attained their present size only after the ice lobes had completely withdrawn from the basins and drainage adjustments had become fully established. Following is a discussion of the ancestors of the Great Lakes in the order of their formation and appearance as the glacier melted northward from the continent.

Lake Maumee

The earliest lake to develop in the Great Lakes area was related to the melting of the ice front of the Erie lobe eastward from the position of the Fort Wayne moraine. This lake developed in the Maumee basin of northwestern Ohio and neighboring parts of Indiana and is referred to as Lake Maumee (Fig. 4). According to Leverett, this lake came into existence about 30,000 years ago. In its early stages, it was confined in a small basin situated between the Fort Wayne moraine and the ice front then standing on the Defiance moraine. The water level of this early basin as shown by shore lines stood at an elevation of 785 feet or 212 feet above the level of present Lake Erie. The drainage outlet was through a narrow valley or col in the moraine which led past Fort Wayne, into the Wabash River and thence south into the Mississippi. A narrow arm of this early lake extended along the border of the ice to the vicinity

of Ypsilanti where a remnant of the shore may be traced along Summit Street at an elevation of approximately 800 feet.

In its early stages, Lake Maumee occupied a relatively small depression, but as the glacier oscillated in retreat the basin was progressively enlarged with the result that the water level was forced to drop as is evidenced also by beach and shore features.

When the ice melted back and uncovered the area on the "Thumb" near Imlay City, the waters of the expanded lake Maumee found a new outlet about 35 feet lower than the Fort Wayne channel and then the waters of the lake flowed out through the "Imlay channel," at an elevation of 750 feet. From the Imlay channel the drainage water was carried west-



Figure 4.—Map Showing Lake Development in the Maumee Stage.
(Modified after Leverett and Taylor).

ward down the Flint River valley past Flint and finally into the Grand River near Lyons. Thence by this drainage way, it flowed into Lake Chicago, the predecessor of Lake Michigan, near the present site of Grand Rapids. This ancient river which drained the ancestor of Lake Erie, is now a valley much too large for the swamp and the small streams now occupying and flowing through it.

During the Imlay outlet stage of Lake Maumee, the ice border rested on the Flint-Mayville and St. Johns moraines. When the ice of the Saginaw lobe finally melted back into the Saginaw basin sufficiently far to expose a land surface with a general slope toward it, lake waters accumulated between the ice and the westward margin of the trough to form early Lake Saginaw. This lake found an outlet through Maple River near Maple Rapids from whence it discharged its waters into the Grand River at Lyons and thus westward across the state to Lake Chicago. Shore features in the early Saginaw lake plain show the Maple River outlet to have stood at an elevation of 730 feet or 20 feet below the Imlay outlet of Lake Maumee.

Lake Arkona

In the next episode of lake history the ice front withdrew altogether from the "Thumb" of Michigan and another lake, Arkona, came into existence. The ice margin retreated to a position some 25 miles north of Bad Axe, the waters of the expanded lake Maumee fell to the level of Lake Saginaw and gradually merged with it. The drainage waters of the lake went around the "Thumb" and found an outlet through the Maple River into the Grand River. The highest beach of Lake Arkona stands at an elevation of 710 feet above sea level at the outlet. Down cutting by the large river which drained lake Arkona through the Grand River channel lowered the water plane to 694 feet. Lake Arkona was of short duration. It was not drained but a readvance of the ice over the "Thumb" separated the waters in the Saginaw and Erie basins, destroyed Lake Arkona and raised the level of the water in the Erie basin to form a larger lake which is named Lake Whittlesey.

Lake Whittlesey

During the Whittlesey stage the ice front maintained a temporary position on the main moraine of the Port Huron system (Fig. 5). The section of the moraine between Vassar in Tuscola County and Bentley in northern Bay County stood in 50 feet of water. At that time the Erie lobe had melted back to the position of Lake Ontario and formed the Erie-Ontario lobe which was separated from the Huron lobe, the ice in the Huron Basin, by an island of exposed land some fifty miles or more across, located in Ontario (See Map, Figs. 4, 5).

The waters of Lake Whittlesey discharged northward through the Black River embayment (the first stage of lower Lake Huron) along the outer border of the Port Huron moraine (now in Sanilac County) and found an outlet near the site of the village of Uby, Huron County. From here the drainage was directed westward through the channel now occu-

pied by the Cass River into independent Lake Saginaw where it eventually found its way out to the Grand River by way of Maple River. The highest beaches of Lake Whittlesey stand at an elevation of 740 feet. Before the close of this stage, however, the Maple River channel was deepened by scouring and the lake level dropped to 735 feet.



Figure 5.—Map Showing Lake Development in the Whittlesey Stage.
(Modified after Leverett and Taylor).

The water of Lake Whittlesey was maintained at the level of the Ubyly outlet only while the ice front rested on the Port Huron moraine. That the ice maintained this position for a relatively long interval, is manifested by the strong beaches and distinct shore features which may be traced for miles along the Lake Huron side of the "Thumb" in eastern Michigan. When the glacier finally retreated up the "Thumb," the lake level dropped so low that no water could discharge through the Ubyly outlet.

When the large lake Whittlesey occupied and overflowed the Erie Basin the ice of the lake in the Lake Michigan trough had melted back far enough to uncover a small basin at its south margin which was filled with water from the melting ice to form Lake Chicago. The ice still filled the Superior basin but had started to melt slightly in the vicinity

of Duluth forming Lake Duluth (Fig. 5). Small fingers of ice extending out from the Ontario lobe in New York, found their way into river valleys which were gradually sculptured into Finger Lakes. These had an independent drainage southward through the Susquehanna River into the Atlantic.

Lake Wayne

Lake Wayne, so named because its beaches are prominent at Wayne, Wayne County, followed Lake Whittlesey. It formed when the ice of the Ontario lobe in the east melted back to uncover a portion of the Mohawk valley. The Uby outlet was abandoned and the drainage of



Figure 6.—Map Showing Lake Development in Warren Stage.
(Modified after Leverett and Taylor).

the lake was directed eastward along the edge of the ice past Syracuse, New York, and into the Mohawk to the Hudson River. During this stage, the Finger Lakes of New York were also connected with the main line of drainage. This is the first occasion in the history of the Great Lakes that the drainage was eastward and the Grand River was not utilized for discharge. A slight readvance of the ice in the east, however, soon blocked the Mohawk outlet and brought the Wayne stage to a close.

Lake Warren

Lake Warren which followed Lake Wayne was formed in the Erie Basin when the ice barrier stood slightly north of Alpena in Michigan and south of Rochester in New York (Fig. 6). Its waters merged with those in the Saginaw basin and gradually expanded to include the water in the Huron-Erie basin as well. The western outlet was again lower than the New York-Mohawk outlet and the discharge from the lake was around the "Thumb", into Lake Saginaw and out through the Maple River. Thence westward by way of the Grand River drainage line to Lake Chicago. The Warren beach stands at 670 to 680 feet above sea level or merely a few feet higher than the outlet except in that area to be discussed later where continental uplift has upcanted the beaches like the lift of a trap door.



Figure. 7.—Map Showing Lake Development in Lundy Stage.
(Modified after Leverett and Taylor).

Lake Lundy

Lake Lundy which followed Lake Warren was relatively short lived and was merely transitional to the succeeding stage, Algonquin (Fig. 7). In the Lundy stage the ice barrier had retreated to a position north of Alpena in Michigan and rested between the Niagara escarpment and

Lake Ontario in the region of Niagara. The outlet was similar to the outlet of Lake Wayne, as the ice in the east had retreated northward sufficiently far to again open the Mohawk outlet past Rome, New York, and so permit drainage in the direction of the Hudson River. Owing to the opening of a lower outlet, the Lundy lake level dropped to between 620 and 640 feet. Later, with the complete withdrawal of the ice sheet from the lowland area south of Ontario, the waters of Lake Lundy were drawn down to a level which uncovered the cliff at Niagara over which the waters spilled and allowed Niagara Falls to come into operation. This is the first evidence of discharge through the Lake Erie to Lake Ontario route. At the Lundy stage in the Erie-Huron Basin Lake Chicago in the Michigan Basin had enlarged northward about to Charlevoix and Lake Duluth filled the western part of the Superior Basin.




Figure 8.—Map showing Lake Development in Algonquin Stage.
(Modified after Leverett and Taylor).

Lake Algonquin

The last of the glacial lakes is known as Lake Algonquin. This stage followed Lundy and represents the greatest expansion of water in the development of the Great Lakes depressions. The great mass of ice had entirely melted from the Great Lakes region. The various lakes in the

different basins were merged into one vast body of water (Fig. 8), Lake Algonquin. In the culmination of this stage, the lake extended northward to include the Nipigon basin in Canada. The eastern lowlands of the northern Peninsula excepting a few small islands were submerged. During this stage the Strait of Mackinac was opened to connect the overflow lakes in the Superior, Huron, and Michigan basins.

As the weight of the ice was removed from the northern part of North America the continent began to rise, to be uptilted along several definite "hinge lines" much as a trap door is upcanted along its hinges. From Algonquin time to the present this upcanting of the continent has caused changes in the Great Lakes. These uplifts in Algonquin time caused several changes in drainage in the Algonquin lake history. An early outlet of Lake Algonquin was opened at Kirkfield, Ontario, soon after the ice had withdrawn from the Superior Basin. The discharge was then carried from Georgian Bay (of the Lake Huron Basin) through the Trent Valley to the Ontario basin and Lake Erie was cut off from the northern lakes. As differential uplifts continued, the Kirkfield outlet was raised so high that it soon became abandoned. For a short interval following the closing of the Kirkfield outlet, the discharge of the lake was divided more or less between lower outlets at Chicago and to the Mississippi and at Port Huron and back to Lake Erie and the Atlantic. At Chicago, a sill of resistant limestone effectively slowed up the process of downcutting. At Port Huron, the drainage waters passed through an area of less resistant clayey material which was readily eroded. Consequently, the Port Huron outlet was soon lowered below the Chicago outlet and the discharge from all the lakes was diverted wholly into the Port Huron line of drainage.

The beaches of Lake Algonquin which must have been level and horizontal when made present striking evidence of the uptilting of the continent. In the area of horizontal beaches south of the hinge line which crosses the state approximately from Port Huron to Manistee, the beaches of higher Lake Algonquin stand at a level of 607 feet above the sea. But at Kirkfield the shore features are at an elevation of 883 feet above sea level which means that in the distance from Port Huron to Kirkfield, the land has been raised 276 feet by differential uplift. Farther north, in the region of Georgian Bay, and the north shore of Lake Superior, the region has been raised at least 600 to 700 feet. In the high morainic hills south of Munising, the highest Algonquin shore is found at an elevation of 948 feet above sea level and a few miles south of Grand Marais, the morainic bluffs show evidences of Algonquin shore work to an elevation of 890 feet. At Grand Marais, however, the successive lower Algonquin beaches can readily be traced and show definitely the progressive changes in lake level which have taken place.

Throughout the different intervals of Lake Algonquin history, variable volumes of water flowed across the Niagara Falls. At first, when the upper lakes used the Kirkfield outlet, the Niagara River carried only the discharge of Lake Erie, which was then about 15 per cent of the present volume. Finally, as the Port Huron outlet assumed control of the drainage, a full volume of flow by this route resulted. The effects of the changes in Lake Algonquin drainage are readily observed in the gorge of the Niagara River in the section between Lewiston, on the original escarpment, and the whirlpool.

Lake Nipissing

When the glaciers had retreated northward well beyond the border of the Superior basin, the land was raised to the point where the Kirkfield outlet was permanently closed. A new and lower outlet was opened at North Bay with discharge into the St. Lawrence River by way of the Ottawa River. Thus was initiated the post-glacial or Nipissing stage of lake history.

The Nipissing Great Lakes occupied the basins of Lakes Huron, Michigan, and Superior. Lake Erie was independent of the upper lakes during the time they discharged through the Kirkfield and Ottawa River outlets. The Nipissing shore is well formed and may be easily traced along beaches which stand at an elevation of 595 feet, above sea level. Continued uplift of the land north of the hinge line eventually raised the North Bay outlet sufficiently high to close it to further discharge. The water was then directed southward into the St. Clair and Port Huron outlets and the present lake stage with its drainage through the Niagara and St. Lawrence Rivers was introduced. In this stage, Michigan attained her present physiographic complexion.

During the interval of maximum glacier invasion, the millions of cubic miles of ice resting upon the surface of the continent exerted a tremendous weight upon the rock formations below. This pressure was sufficient to cause depression of the entire Laurentian plateau with an ultimate downwarping estimated at 800 to 900 feet in the region north of Lake Superior. The downwarping of the continental block allowed salt water of the Atlantic Ocean to invade the Hudson and St. Lawrence Valleys and drown the basin of Lake Champlain and the upper end of Lake Ontario. This inundation is commonly referred to as the Champlain Marine stage. A series of clays was deposited in the waters during this stage and in the clays the remains of a few animals which could exist in cold sea water, a sub-arctic fauna, are found. The partial skeletons of two whales recently discovered by Dr. E. C. Case of the University of Michigan, in the bogs overlying glacial deposits along the

Huron shore in Michigan suggest a possible open route from the ocean to the fresh water lakes through which the whales migrated inland, but does not imply that the Great Lakes were once bodies of salt water.

The final removal of the vast sheet of continental ice and the consequent release of pressure due to melting has allowed the depressed region to partially recover its original attitude. (Fig. 10). Differential uplift, which began possibly with the first recession of ice in the Wisconsin stage, is still in progress—a fact which seems to bear out the conclusion that the recovery has not been completed. At present, the




Figure 9.—Map Showing Lake Development in Nipissing Stage.
(Modified after Leverett and Taylor).

land is rising at the rate of 10 inches per hundred years, or one-tenth of an inch per year for each hundred miles of distance north of the Algonquin hinge line. (The line extends as stated earlier from near Port Huron across the State to Manistee.)

The effect of the continued uplifts has been to tilt, and in some places to split, the beaches of the Algonquin and Nipissing Lakes. The amount of tilting to which Michigan has been subjected since the Algonquin Lake stage may be readily measured by comparing the present elevation of its beaches in various localities. In the area of horizontality, where




Figure 10.—Map showing position of hinge lines of uplift and present Great Lakes. (After Leverett).

no uplift has taken place, near Manistee on Lake Michigan, the highest Algonquin shore stands at 607 feet. Six miles north of the Canadian Soo, the same shore is found at 1015 feet. At Port Huron, on the east side of the state, the highest Algonquin beach stands at an elevation of 606 feet. On Mackinac Island, in the Straits, the same beach rises to 809 feet.

The Nipissing beaches likewise show the effects of differential uplift, but on a much smaller scale. In the area of horizontality near Manistee, the Nipissing shore has an elevation of 595 feet. At Sault Ste. Marie, the same shore stands at 651 feet—which evidences an uplift of 56 feet in a distance of 160 miles. At Port Huron, the elevation of the Nipissing beach is 595 feet, but at Cheboygan it is 31 feet higher, or 626 feet.

The main portion of this paper reports on an area which is within the region of the Algonquin and Nipissing Great Lakes.

SYNOPSIS


The Belted Lowland province of the northern Peninsula (Fig. 11), extending from the meridian of Marquette eastward, presents a variety of complex problems in the light of Pleistocene glaciation. This area was the locus of development of the Green Bay and Michigan lobes during the maximum extension of the late Wisconsin ice sheet. Later it became the center of renewed activity in the readvance of the independent lobe from the Lake Superior basin shortly before its final withdrawal from the surface of Michigan.

When the ice of the late Wisconsin stage finally withdrew to the north far enough to uncover the Superior region, the lowlands were inundated almost completely by the waters of Lake Algonquin. A very small proportion of the province had a relief sufficiently high to extend above this water plane. Consequently, the surface deposits left upon retreat of the late Wisconsin ice sheet were, in large measure, influenced by water and wave activity, with the result that they were subjected to widespread and differential modifications.

Moraines which were covered by the lake waters have lost, to a large degree, the characters and features possessed when laid down by the waning ice sheet. The crests were smoothed down, flattened out and in many places, reduced to level plains. Slopes of the stronger systems were washed down, subdued and subsequently became mantled with a thin veneer of lacustrine sand, which locally obscures the original till. In those areas where the moraines have lost their identity as definite features of relief, it is still possible to determine the sites upon which they originally stood, by the presence of till remnants and concentrated erratics; except where the modifying cover has developed to such a depth as to completely mask the former surface.

Post Algonquin drainage ways developed in the region, have successively cut through the morainic systems and dissected them to such a degree that they now stand as dismembered units, often widely separated by low-lying swampy and marshy tracts. The disconnected and water-modified moraines are thus difficult to trace and much of the field evidence, necessary to a satisfactory and complete interpretation of the glacial problems involved, is entirely lacking.

The successive water levels, developed along the prograding shores of glacial Lake Algonquin, are marked, either by definite wave-cut features where the water level was held up for a comparatively long interval, or, by aeolian shore ridges where retreat was more rapid. Many of the retreatal halts in shore recession were centered on morainal plains and



slopes. Here the primary relief became masked by shallow deposits of lacustrine sand which in turn was spread out locally in the form of low, wind-blown waves and ridges of variable extent.

The relief of the Belted Lowlands in the eastern half of the northern Peninsula is relatively quite low. The elevation, in general, does not exceed 250 feet above the Great Lakes, except in restricted areas where it may rise considerably above this level. The low character of the topography is reflected in the extensive tracts of poorly drained marshy, sandy plains and muskeg which abound in this region.

The largest and most continuous area of swampy lowland in the northern Peninsula lies in the drainage ways of the Tahquamenon and Manistique Rivers, confined largely within the borders of Alger, Luce and Schoolcraft counties. This region presents features which are characteristic of the Belted Lowlands as a whole, and unquestionably holds the key to a proper understanding of the problems involved in the Pleistocene glaciation of the province. The drainage basins of the Tahquamenon and Manistique Rivers lie almost wholly within the region which was invaded by a readvance of the Superior lobe of the Labrador ice sheet, following the break-up of the Michigan lobe, late in the Wisconsin stage of Pleistocene glaciation. This lobe moved from the northeast and followed a course which was directed largely by the pre-glacial depression of Lake Superior. The general direction of the ice movement was to the southwest, but it is evident from glacial striae in the eastern portion of the Peninsula, that there was also a southward trend, which was responsible for much of the glacial activity in this area.

The work of the glacier in this region was performed largely in the receding hemicycle of glaciation as stated by Hobbs (1). This is evidenced by the fact that previous to the readvance of the Superior lobe, tongues of ice representing the Green Bay, Michigan, and Huron lobes, which formerly deployed over vast areas in southern Wisconsin and Michigan, had withdrawn from their basins and become broken up completely.

In its readvance, the ice sheet removed much of the older residual soil and weathered rock waste which had accumulated through protracted periods of erosional activity. The bed rock surfaces were planed and scored and large blocks, frozen to the sole of the ice, were plucked out and moved with the glacier. Old valleys, which had been chiseled out of solid rock by pre-glacial streams and rivers, afforded natural thoroughfares for the movement of the ice tongues. They were consequently deepened and widened by the plucking and gouging activity of the deploying ice mass; the higher ridges and exposed rock areas were smoothed down to a large degree and the features in general suffered

a decided reduction in relief. In the process of deglaciation which was induced by recession of the ice sheet, due to dissipation of its marginal areas, there was spread out over the rock pavement a mantle of glacial drift of variable thickness and composition. The pre-glacial valleys were filled quite completely with debris deposited by the waning ice.

The continued wasting of the glacier caused the ice front to retreat farther and farther to the northeast and allowed a slow uncovering of the basin of Lake Superior. The water from the melting ice, collecting in the growing basin, gradually filled it to overflowing and initiated glacial Lake Algonquin, a predecessor to Lakes Superior, Michigan and Huron. In the highest stage, this lake spread its waters over the greater portion of the Tahquamenon-Manistique drainage area, and was responsible for a multitude of changes in the surface expression of the region.

The complete withdrawal of the ice from the east end of the Superior basin opened up the Kirkfield outlet in Ontario and allowed the waters of Lake Algonquin to drain through the Trent Valley. The outlet past Port Huron was likewise functioning during this stage of lake history with the result that drainage was affected both to the east and to the south. The lowering of the Algonquin lake level eventually caused the waters to split along the divide between the present northward flowing Tahquamenon and southward flowing Manistique drainage ways. The parting of the Algonquin waters is definitely evidenced by the complete reversal of slopes in the east-west trending dune ridges on either side of the divide.

The final dissipation of the ice sheet in the Lake Superior basin resulted in a decided reduction of ice weight on the land surface. Differential uplift and tilting, as described by Gilbert (2), Goldthwait (3), Fairchild (4), and Spencer (5), raised the superior region several hundred feet above the normal plane. Taylor (6) ascribes the uplifting movements as due to resilience following depression by ice weight and also to crustal creep. This change in elevation was responsible for decided modifications in drainage and resulted in the development of broad, swampy, and poorly drained tracts along the borders of north-flowing streams. The southward flowing rivers, on the other hand, were simultaneously increased in gradient and, due to rejuvenated erosional power, proceeded to excavate their channels more deeply into the sandy lake floor.

The initiation of fluvial activity on the newly emerged land surface, coupled with the successive lowering of the water level in the slowly receding lake, brought on a new set of geologic conditions. Aeolian activity became a prominent factor and translocation by the wind of sand from the river flats, newly formed lake beaches and terraces, resulted in the piling up of innumerable sand dunes and ridges. These in turn

caused diversion and blocking of river courses, with the ultimate result that the drainage systems became completely deranged and extensive swampy and marshy tracts developed. Thus, the surface features became woven into an intricate and extremely complicated pattern of undifferentiated complexes, at times very difficult to interpret.

INTRODUCTION

The field work, which represents the basis for this report, was pursued during the field seasons of 1928, 1929, 1931, and 1932. The work was carried on in connection with the Land Economic Survey which was attempting to obtain a complete inventory of conditions in certain northern counties.

The territory under consideration is, for the most part, a heavily timbered wilderness and facilities for travel are in places quite limited. All of the passable roads as well as many old railroad grades, grown over logging trails and abandoned tote roads, were traveled by automobile. Hundreds of miles of trails were followed on foot and much of the inaccessible area was covered by the ordinary pace and compass method.

In the field work of Schoolcraft County covered during part of the season of 1931 and all of 1932, a complete set of aerial photographs which were of considerable aid in working out the many details of surface structure and tying the features in to the general land net, were at the writer's disposal. These photographs prepared by the aerial parties of the United States Geological Survey were later assembled into mosaics by the State Geological Survey for use in the field.

The problem herein discussed suggested itself in connection with the complications involved as the field work progressed. The area in question seemed to contain the key to the interpretation of the surface geology of the low swampy areas which constitute such a large portion of the Lowlands province in the eastern half of the Northern Peninsula.

Known elevations were obtained from profile maps of established railroad grades. These served as the basis for running elevations by means of aneroids. The aneroid readings were checked and rechecked on the established known points and corrected each day from the records of the barograph which was located in the main camp.

ACKNOWLEDGMENTS

The author is greatly indebted to Professor I. D. Scott whose visit into the field during the summer of 1930 prepared the way for the solution of some of the intricate problems involved in the aeolian geology of the Manistique drainage basin. Also, for many valuable suggestions received from him during the progress of the work. Dr. Frank Leverett has checked the manuscript and maps and otherwise offered expert opinions concerning certain areas which seemed in the field to have very obscure geological relationships. The author is grateful to Dr. W. H.

Hobbs for suggestions and criticisms which have been offered from time to time and for his constructive interest in the problem. Dr. R. A. Smith, State Geologist, has made valuable contributions concerning field methods, and through the course of the work has shown keen interest in the developments. It is largely through his efforts that it was possible to carry the field work to completion. In field and office, Mr. L. R. Schoenmann, of the Land Economic Survey, has imparted important information concerning many features of the problem as related to soil classification. His keen observations in the field, together with the sound interpretations always made of them, have been of utmost value in checking personal conclusions.

The author wishes to express thanks to the directors of the State Geological Survey and the Land Economic Survey for the use of field equipment of all descriptions, maps and photographs which were always made available. No small amount of credit is due the men of the Land Economic Survey field crews who through the four seasons of field work have contributed much information concerning corners, boundaries, swamps, lakes, and a multitude of other features. Space does not permit the listing of the names of the many individuals who have thus shown an interest. Mr. O. F. Poindexter, of the State Geological Survey, has offered valuable suggestions in the field, and also made checks on elevations in various parts of Schoolcraft County. The drafting of the final maps from the field sheets has been the work of Allerd W. Bergquist, to whom the writer desires to express appreciation. To Miss Helen Martin who has so painstakingly edited the volume, the author is greatly indebted.

Chapter I

THE TAHQUAMENON DRAINAGE SYSTEM

GENERAL CONSIDERATIONS

Approximately 40 per cent of the total area of 910 square miles in Luce County is covered with poorly drained swamps, marshes, and low-lying sand plains. The greater portion of this region lies within the Tahquamenon drainage basin.

The Tahquamenon River rises in a series of glacial pit lakes in T. 47 N., R. 12 W., in Luce County, at an altitude of approximately 800 feet above seal level or 200 feet above Lake Superior. For a distance of ten miles from its source, the river flows generally southward, through an extensive outwash plain, a few miles east of the county line. Near the center of T. 46 N., R. 12 W., the river swings abruptly eastward, into the swampy depression, and continues in that direction across the county. Thence it turns northward, and leads out of the county in the northeast corner of T. 48 N., R. 8 W. Its waters are finally discharged into Whitefish Bay in Lake Superior, near Emerson, Chippewa County.

For much of its course in Luce County, the river flows through an extensive, poorly drained lowland, composed largely of organic matter resting upon lacustrine sand and underlain at varying depths with a reddish-colored, plastic lake clay. Many tributary streams, most of them small, gather their headwaters at the margin of this lowland and at times contribute considerable volume to the main river. Some of the larger streams of this system are fairly well developed and reach back for considerable distance beyond the swamp. The broad interstream areas are low, swampy, and poorly drained. The drainage pattern is not at all systematic, but rather, extremely haphazard in development.

From the point where its course turns eastward in T. 46 N., R. 12 W., and for a distance of fifty miles, to the Upper Falls, the river descends 26 feet. Eight feet of this fall is concentrated in a distance of three-fourths of a mile above the falls and probably represents the amount of down cutting since the rock barrier was cut through. The average gradient of the river above the Upper Falls is less than 6 inches to the mile. The channel is shallow and poorly confined, and in the periods of heavy rains and melting snows does not begin to accommodate the vast volume of water which is brought to it by the tributaries from the water soaked, spongy lowland tracts. Consequently, during the flood seasons the water overflows the channel banks, backs up into the drainage ways

PLATE I



Figure 1.—Tahquamenon River above the Upper Falls.



Figure 2.—The Upper Falls of the Tahquamenon River.

and spreads out over the low interfluves to form extended shallow lakes. (Plate I, Fig. 1)

At the Upper Falls, (Plate I, Fig. 2,) in Section 11, T. 48 N., R. 8 W., the river plunges over a forty foot ledge of Cambrian sandstone and for some distance below is carried through a narrow gorge which has been sculptured out of the same material (Plate II, Fig. 1). In this area the gradient is much steeper, being approximately 7 feet to the mile, in the distance of four miles between the base of the Upper and the crest of the Lower Falls. Its velocity in the gorge is very great and its cutting power at a maximum. The elevation of the crest of the Upper Falls, as determined by Wisler (7), by means of levels, is 694 feet. The Lower Falls in Chippewa County stand with the crest at an elevation of 626 feet and drop 20 feet to a level just 4 feet above the water plane of Lake Superior.

The watershed between the Tahquamenon and Manistique drainage systems lies near the line between Luce and Schoolcraft counties, a short distance north of Danaher, on the Duluth South Shore and Atlantic Railroad. The divide is in the swamp at an altitude of 720 feet, or approximately 120 feet above the level of Lake Superior.

Late in Algonquin time, the Tahquamenon depression provided a drainage connection between the Superior and Lake Michigan basins. The discharge waters from the upper basin were carried through the narrow strait, north of McMillan, across the divide and south into the Manistique River system. This outlet remained open until the level of Lake Algonquin was depressed to a plane below the level of the divide. At this stage in late Algonquin time, the two drainage systems were separated; the Tahquamenon waters were diverted into Lake Superior and the Manistique drainage leading southward into Lake Michigan.

The early development of the Tahquamenon basin is associated with geologic activity which antedates the glacial epoch. The present valley occupies in part an old broad, pre-glacial trough, (Plate II, Fig. 2), which had been excavated quite deeply into bed rock before it was invaded by Pleistocene glaciation. The old depression was fairly well filled by deposits left as the ice retreated and also by subsequent alluviation. It became the site for the present river shortly before the waters of the Superior basin had reached the Nipissing stage.

The southern border of the Tahquamenon basin is confined by the steep, northward facing slope of the Niagara cuesta. This escarpment, which forms the south wall of the basin in Luce County, (Fig. 12) has played a very important role in the geologic development of the eastern half of the Northern Peninsula. It forms the backbone structure of the section and is also the axis of Green Bay peninsula in Wisconsin. The Tahquamenon depression bears the same general relationship to the escarpment

PLATE II



Figure 1.—Sandstone Gorge below the Upper Falls of the Tahquamenon River.




Figure 2.—Looking north across the Tahquamenon Basin from the crest of the Niagaran Escarpment, just north of Hendricks Quarry.

MICHIGAN
DEPARTMENT OF CONSERVATION
LAND ECONOMIC AND GEOLOGICAL SURVEYS
COOPERATING

→

GEOLOGICAL MAP
SHOWING
SURFACE FORMATIONS
OF
LUCE COUNTY





KEY TO MAP

SURFACE FORMATIONS

- SAND DUNES
- SANDY LAKE PLAINS
- CLAYEY LAKE PLAINS
- OLD SHORE LINES
- ALGONQUIN
- UPRISING
- INDEFINITE

- MORAINES ABOVE THE WATERS OF GLACIAL LAKES
- MORAINES COVERED BY WATER
- MORAINES OF GLACIAL LAKES
- OUTWASH PLAINS
- ESCARPMENTS
- SWAMPS PEAT
- MARL

BASE FEATURES

- CITIES
- VILLAGES
- CAMP
- RAILROADS
- ROADS
- RAILROADS
- GRAVEL PITS
- ELEVATIONS
- BAROMETRIC
- QUARRIES

in Luce County as does the Green Bay depression to its adjoining peninsula farther south. In fact, the two depressions are somewhat continuous and possess a similar geologic origin and general structure.

The differential uplifts, which strongly affected the region during later Algonquin time, raised the rock ledge at the Upper Falls and caused a decided reduction in the upstream gradient of the river. The velocity was decreased to such an extent that degradational processes were largely terminated. The waters of the river became impounded behind the slowly rising rock barrier and were forced to spread out over the bordering lowland tracts for considerable distances back from the channel. An extensive shallow lake, thus formed in the drainage basin of the river, became the locus for sedimentation of material carried in from the higher uplands by various tributaries. Through the ingrowth of hydrophytic vegetation, the lake gradually assumed a swampy condition and became the site for successive accumulations of peat and muck.

Throughout the major swampy lowlands of the Tahquamenon and other drainage areas, an intricate system of sandy ridges and narrow plains has developed. The ridges range in size from a score of feet, to several miles in length, and from a few feet, to twenty and more feet in height. They are composed essentially of subangular grains of quartz sufficiently small to be easily transported by the wind. The ridges are not always uniform in composition and often grade into coarse sand and gravel towards their bases. In many cases they are coarse textured throughout the entire structure.

The general trend and arrangement of the ridges, together with the texture of the material contained in them, suggests an aeolian origin. Many of the ridges possess the contrasted leeward and windward slopes so characteristic of dunes. The gravelly sections of the ridges, however, are of lacustrine deposition and were probably formed as off-shore deposits along the margin of the receding Lake Algonquin. It seems logical to assume then, that the numerous ridges have had a two-fold origin and were formed by the wind blowing the finer sand up into definite features from the old water laid beaches. The transverse dunes which cross the Tahquamenon valley have their steep, leeward slopes directed universally to the south, a feature which would suggest a northward retreat of the Algonquin waters.

Along the borders of the swamps, and especially where they merge with the higher glacial features, sandy tracts have been shaped into relatively smooth, level plains. These are sometimes perched rather high up on the slopes of moraines and outwash plains, up to elevations of 750 to 800 feet. McMillan and Newberry, situated on the inner border of the outer morainic system, are both located on such a sandy plain.

The sandy ridges and narrow plains which ramify through the swampy

areas, are generally covered with a growth of pine and other vegetation. They afford more or less connected thoroughfares of travel for both man and animal, through an otherwise unbroken jungle. They also provide sites for camps and shelters for hunters and trappers who invade these areas. The ridges furnish fairly suitable material for the construction and maintenance of such roads as penetrate the wilderness in this region.

In general, the areas of muck are so low and poorly drained that they cannot be utilized for agricultural purposes. Some of the clayey deposits, especially those which are high and fairly well drained, have been planted to alfalfa and clover with good results. These heavier soils, however, retain so much moisture and remain cold for such longer periods in the spring, that early planting is very difficult.

Several other drainage systems, in no way allied to the present Tahquamenon basin, but, nevertheless, important because of their association with lowland areas in Luce County, will be given brief consideration in this text. They include in order of importance:

- A. The Two Hearted River System.
- B. The Little Two Hearted River.
- C. The Sucker Basin.
- D. The Shelldrake Drainage System.

A. THE TWO HEARTED RIVER SYSTEM

The Two Hearted River has its headwaters in a series of swamps in a region north of the Munising morainic system in T. 48 N., Rs. 11 and 12 W. The river follows a northeasterly course through tracts of low swamps and higher sandy lake plains, and discharges into Lake Superior in T. 50 N., R. 9 W. This river system drains an area of 230 square miles, all in Luce County, and maintains a rather constant flow throughout the year. In its upper reaches, the river leads through an extensive swampy lowland. In the lower half of its course, through the old lake plain, it has incised its channel quite deeply into loose sand and is lined with rather steep banks. For a distance of about five miles back from its mouth, the river almost parallels the shore of Lake Superior and is prevented from directly entering the lake by a row of low sand dunes which fringe the shore.

The Two Hearted drainage system, like that of the Tahquamenon River, was greatly disturbed by the differential uplifts which affected the region. The gradient of this river was likewise reduced and the velocity and erosive power retarded. Unlike the Tahquamenon River, however, there was no rock threshold near its outlet to hold the waters back. Consequently, the river, able to deepen its channel, has not suffered the various relapses which mark the history of the Tahquamenon River.

B. THE LITTLE TWO HEARTED RIVER.

This river has its source in the group of Two Hearted Lakes in T. 48 N., R. 9 W., and drains an area of approximately 75 square miles in the northeastern part of the county. It is a more or less independent river and has relatively few tributaries and branches. The flow is generally northward, across a low, sandy lake plain and it empties into Lake Superior in T. 50 N., R. 9 W., a few miles east of the outlet of the Two Hearted River.

From near its headwaters, the river swings with a broad bend in a northeasterly direction around a group of lakes which are concentrated in the S. E. $\frac{1}{4}$ of T. 49 N., R. 9 W. These lakes apparently occupy the site of a former morainic surface, probably a continuation of the Munising morainic system which has been eroded down, possibly by stream as well as by lake action.

C. THE SUCKER BASIN.

The Sucker Basin, in the northwestern corner of Luce County, carries the waters of the Blind and Dead Sucker rivers and drains an area of 45 square miles in the County. These two rivers have very low gradients, are somewhat stagnant in flow and bordered by broad areas of low swamp. They are separated from each other by a low, sandy interfluvium of sufficient elevation to have held in the waters of Lake Nipissing. No definite shore line features are traceable along this divide, however; all evidence having been erased by subsequent water action.

The south rim of the basin is marked by a steep, wave-cut escarpment, with a crest at an altitude of 730 to 750 feet. This wall served to hold in the waters of the receding Lake Algonquin as it reached its lower stages. The north edge of the basin is hemmed in by a series of dunes and sandy ridges which rise to a height of 75 to 100 feet and more above the level of Lake Superior.

The basin has been subjected to a considerable amount of scour by water action during its occupation by lakes, and still more recently by the rivers which have occupied the basin since glacial times. Differential uplifts in the region have been largely responsible for the reduction of the gradient of the rivers and also for the development of the extensive mucky areas which are enclosed by the basin.

D. THE SHELLDRAKE DRAINAGE SYSTEM.

In the northeastern corner of Luce County is an area of about 24 square miles which lies in the headwaters of the northeastward flowing Sheldrake River. This river is an independent one and has the bulk of its

water-shed in the adjoining areas of Chippewa County. The greater portion of the Shelldrake drainage system in Luce County is in the region of Betsy Lake where it taps a broad, swampy, fringing lowland area.

LAKES

Several hundred lakes, most of them small and many merely ponds, are scattered through the county. They reflect an origin which is intimately associated with the activities of the glacier and lie within areas of moraines, outwash plains and lake plains. Very few lakes are found in the swamps along the major drainage ways.

Lakes which have been formed in the morainic areas are referred to as *morainal* types. They occupy depressions set in between the knolls and ridges of the more rugged topography. They are generally bordered by upland slopes which may afford favorable sites for resort development. Manistique and North Manistique lakes are the largest basins of this character; others of smaller size are widely distributed through the several morainic systems. Lake basins of this type were formed either by the gradual melting of detached blocks of glacier ice which had become buried under glacial debris, or through a process of "push and dump" by the glacier. Many of the depressions not occupied by water, were likewise formed in this manner and give character to the swell and sag expression of the surface.

Lakes which occupy the outwash and lake plain areas are generally referred to as *pit lakes*. They have a geological origin similar to that of the kettles in the moraines and often are so numerous as to give to the surface a pitted relief. Pit lakes are commonly bordered by level tracts of sandy material and are usually not so deeply depressed as are the basins in the moraines. Several large depressions of this type are located in the level plain to the north, among them being Muscalonge, Bodi, Temple and Pike lakes. Lakes of this character have a fairly good bottom and shore except where their margins have been encroached on by the ingrowth of vegetation and the deposition of peat.

Occasionally, lakes were developed with basins partially set in between morainic ridges and adjoining outwash plains. Such forms are classed as *fosse* lakes. They differ from the other types merely in position with reference to surface features; geologically, they have a similar origin. Some of the Tahquamenon lakes along the west border of the middle morainic system and others such as Murray and Goose lakes farther east present this feature.

Lakes which are situated in the drainage ways are generally bordered with low swampy tracts which make them almost wholly inaccessible.

Mud Lake, north and east of McMillan, and lake Betsy in the northeastern part of the county, are good examples.

The lakes in Luce County have a potential recreational value which is rather difficult to evaluate. They are visited each summer by hundreds of tourists who enjoy their settings in the timbered wilds. One of the chief obstacles in the way of extensive resort development among these lakes is the lack of facilities for getting to them. Roads, in certain parts of the county are entirely lacking and many of the lakes may be reached only after hours of travel over undeveloped trails, which stretch through the wilderness. With a good system of passable roads and a definite conservation policy of stocking the lakes with fish, it is quite certain that many such isolated areas would furnish playground and recreational facilities for many thousands of tourists who now are not familiar with the region.

SWAMPS AND MARSHY TRACTS

Swamps, marshes, and muskeg cover about 347.62 square miles or approximately 39 per cent of the surface in Luce County, with a distribution in the various townships as indicated in the table below.


<i>Township</i>	<i>Square Miles</i>	<i>Township</i>	<i>Square Miles</i>
T. 45 N., R. 8 W.	19.58	T. 48 N., R. 8 W.	9.62
T. 45 N., R. 9 W.	12.75	T. 48 N., R. 9 W.	12.05
T. 45 N., R. 10 W.	7.09	T. 48 N., R. 10 W.	22.40
T. 45 N., R. 11 W.	6.81	T. 48 N., R. 11 W.	23.36
T. 45 N., R. 12 W.	16.81	T. 48 N., R. 12 W.	11.08
T. 46 N., R. 8 W.	22.45	T. 49 N., R. 8 W.	18.19
T. 46 N., R. 9 W.	22.63	T. 49 N., R. 9 W.	5.04
T. 46 N., R. 10 W.	20.68	T. 49 N., R. 10 W.	2.57
T. 46 N., R. 11 W.	23.28	T. 49 N., R. 11 W.	.96
T. 46 N., R. 12 W.	16.87	T. 49 N., R. 12 W.	9.24
T. 47 N., R. 8 W.	10.44	T. 50 N., R. 8 W.	1.97
T. 47 N., R. 9 W.	26.41	T. 50 N., R. 9 W.	—
T. 47 N., R. 10 W.	22.71	T. 50 N., R. 12 W.	—
T. 47 N., R. 11 W.	2.05		
T. 47 N., R. 12 S.	0.58		

Some of the swampy areas are filled-in lakes of rather small dimension but generally the swamps occupy regions of underdeveloped drainage and are restricted to the low lying flats along the river systems. The swamps are unusually level and for the most part covered with shallow deposits of peat and decaying vegetation. Clay of a very tight and impervious character underlies the poorly consolidated peat in much of the lowland

tract, and is responsible to a large extent for the high position of the water table and the soggy condition of the surface.

The higher outer borders of the swampy lands are generally quite heavily wooded and give growth to pole-size and often log-size trees of tamarack, cedar and spruce. The lower inner reaches, along the stream margins, on the other hand, are quite generally grown up to a tangle of alder and other low brush which makes travel exceedingly difficult. Some of the swamps are quite open and may well be classed as muskeg. Blueberries seem to find a desirable habitat in the drier open bogs and furnish a reasonable source of income to a fairly large number of people during the short season in the late summer. Many of the more poorly drained lakes and basins are margined with quaking bogs and are thus inaccessible. The drainage lines in the lowlands are not sufficiently developed to carry off the surplus water during protracted periods of heavy rains and melting snows, with the result that a large portion of the area cannot be adapted to agricultural purposes. In view of the impossibility of developing much of the land for the production of farm products, it seems that its future value lies principally in the maintenance of a forest and ground cover for the protection and feeding of wild life and the growing of timber.

The swamps have a very high capacity for absorbing water and serve as reservoirs to regulate the flow into the rivers, thus preventing excessive floods which might otherwise occur.



trac
wat
T)
heav
tam:
mar:
alde
of tl
berr
a rea
shor
and
The
off t
ing
adap
ing r
futu
cove:
timb
Th
as re
sive

Chapter II

THE MANISTIQUE DRAINAGE SYSTEM

MANISTIQUE RIVER

Manistique River, which drains the greater portion of Schoolcraft County (Fig. 13), is one of the largest streams in the northern Peninsula. It has a drainage area of about 1,400 square miles, situated mostly in the great Manistique swamp, of which it forms the eastern margin. The main stream follows a general southwesterly direction and enters Lake Michigan at the city of Manistique.

Numerous parallel tributaries, with headwaters occasionally reaching back into the Munising Moraine, within a few miles south of the Lake Superior shore, flow southeastward across the sandy, marshy lowland and enter the trunk stream almost at right angles. These flow with fairly good gradients down the warped slope of the low, marshy lake plain and supply a large volume of water to the main river.

Several large lakes, among the largest in the Northern Peninsula, are tributary to the Manistique River and contribute appreciably to its flow. The most noteworthy of these lakes, the Manistique, with an area of 15.8 square miles, the Whitefish with 6.4 square miles, and the Indian with an area of 13 square miles, have been described in detail by Scott (8).

The main course of the Manistique, from its headwater confluence with the East Branch of Fox River and Fox River, (Plate III, Figs. 1 and 2), follows closely along the eastern edge of the extensive marshy lowland. Throughout much of its distance, the river flows through a poorly drained, marshy, lake plain, and has trenched a fairly well defined channel into the loosely consolidated, spongy floor. In its more youthful upper reaches, where the gradient is higher and the velocity more rapid, the stream has succeeded in maintaining a relatively straight course. In its older, lower stretches, on the other hand, the river braids across the marshy, sandy floor in a series of broad meanders. Many of the larger bends have been entirely cut off and stand as oxbow lagoons, filled with water. The condition of stagnation which was responsible for the development of the meanders and cut-offs, to such a large extent in the lower course of the river, may be traced back to Nipissing times. Miss Stevenson (9) mentions the fact that the Nipissing shore which extends as a spit nearly to the outskirts of Manistique, turned the Manistique River westward. The development on this spit, of a row of dunes, piled up to a height of approximately 100 feet above Lake Michigan has been responsible, to a large extent, for the maintenance of this course from Nipissing

PLATE III



Figure 1.—East Branch of the Fox River, one of the headwaters of the Manistique River, in a muskeg swamp. Sec. 33, T. 45 N., R. 12 W.



Figure 2.—The Fox River, near its headwaters, valley incised in a sandy outwash plain. Sec. 21, T. 47 N., R. 14 W.

times to the present. That the water was ponded behind the Nipissing spit is evidenced by the extensive stagnant slough which extends westward from the shore cliff.

The Algonquin uplifts, which were responsible for the damming of the north-flowing Tahquamenon, had a counter effect upon the Manistique and other south flowing rivers. The headwater reaches of this large river system were more greatly affected by the differential movements and were thus raised correspondingly higher than were the lower stretches. This caused the gradient and likewise the velocity to become increased, with the result that downcutting, rather than widespread flooding, was initiated. In its distance of 68 miles from source to mouth, the Manistique River has a total fall of 115 feet. This represents an average gradient of 20 inches per mile as compared with 6 inches per mile for the Tahquamenon above the upper falls.

The numerous tributary streams of the Manistique River flow in a southerly direction across the tilted floor of glacial lake Algonquin. They are roughly parallel in their courses and, like the master river, are bordered by extended tracts of low lying poorly drained marshes and muskeg beset with wind-blown ridges of sand.

Several of the tributaries, including the East and West Branches of Fox River, Driggs Creek, Marsh Creek and the various headwater limbs of the West Branch of the Manistique River, have extended their valleys back to the outer slope of the Munising moraine, in Alger County. Many of these streams flow for a distance of 25 to 30 miles across country before finally joining the Manistique. They drain the numerous morainal and pit lakes which are distributed through much of the upland watershed.

As has previously been inferred, the Manistique River did not come into existence as an independent drainage way until the Lake Algonquin waters had receded to about the level of the Nipissing stage. The Newberry morainic system, which now borders the swamp in fragmentary and disconnected units, was at that time no doubt a continuous feature and served as a barrier to the slope-directed flow of the various tributaries. Consequently, the massing of the drainage water against the inner slope of the moraine, resulted in the development of the major drainage line. It is quite probable that the south end of the Hiawatha segment of the moraine temporarily dammed the waters of the river and caused a ponding behind it. The large volume contributed by the tributaries soon broke through the obstruction and found an outlet into Lake Nipissing, whose margin stood close to the outer border of the moraine in this locality.

Chapter III

GLACIAL ACTIVITY

During the time of greatest advance of the Wisconsin ice sheet, a number of individual ice tongues moved southward from the basin of Lake Superior through pre-glacial valleys which had been established by stream sculpture in the older rock surface. Two of the early lobes, the Green Bay and the Michigan, deployed across the area discussed in this report and were responsible for the development of certain glacial features which have, to some degree, been left unmodified.

FEATURES DEVELOPED BY THE GREEN BAY LOBE

The Trenary Till Plain

Position Within the Axis of the Green Bay Lobe

The till plain which occupies the southwestern part of Alger County, extending from near Eben Junction and southward through Trenary, represents but a small portion of a much larger plain which may be traced westward for a distance of six to ten miles into Marquette County, and to the Ford River, some twenty-seven miles south, in Delta County.

In eastern Marquette County, the plain is limited on the west by a border moraine which extends from near the vicinity of Marquette in a general southerly direction to the village of Bark River in T. 38 N., R. 24 W., Delta County.

On the east, the till plain is bordered by the Au Train—Whitefish drainage depression, within which lie broken fragments of a moraine, flanking in part, both east and west banks of the channel. In this locality the original moraine was dismembered by the scouring action of the river which temporarily carried the drainage from the Lake Superior basin to Little Bay de Noc in Lake Michigan during the early part of Lake Algonquin times. It is now difficult to trace the complete morainal system to its former southernmost extremity. Field evidence, however, seems to point out that the disconnected and fragmentary eastern morainic limb may at some time have connected with the western limb, and formed a sinuous and more or less continuous loop around the southern end of the till plain. Thus, this moraine marks the borders of the ice in the Green Bay lobe in its final halt during retreat, and emphasizes a definite demarcation between the drumlin-till plain area of Menominee County to the south, and the Trenary esker-till plain area to the north.


PLATE IV



Figure 1.—Portion of Trenary till plain to the north of Trenary.
Esker ridge in background.



Figure 2.—Esker ridge on Trenary till plain, near Limestone.



The ice lobe, which was responsible for the development of the Trenary till plain, pushed through the low depression west of Munising, in Alger County. Its final extent and position are marked definitely by the location of the associated marginal morainal fragments which skirt the plain. The direction of the ice in the lobe was mainly south or west of south on the east side of Big Bay de Noc as evidenced by scattered glacial striae on the exposed rock pavement located by Leverett.

Topography and Relief

The Trenary till plain in Alger County (Plate IV, Fig. 1) stands at an elevation ranging from almost 1,100 feet in the vicinity of Dorsey in the north, to 835 feet in the area southeast of Trenary, and extends well above the highest water plane of Lake Algonquin. The surface slope is generally from northwest to southeast with most of the drainage channels thus controlled. The plain as a whole is generally quite smooth but locally assumes a somewhat wavy to undulating character. In places, where the drift is thin, the surface expression is influenced directly by the character of the underlying rock surface. Eskers are numerous and widely distributed throughout the extent of the plain. They are extremely variable in dimensions, ranging from merely a few hundred feet to several miles in length and from ten to fifty feet in height. They are generally narrow, although where several coalesce they may be expanded to form broken features a half mile or more wide.

A few drumlins in the vicinity of Eben Junction, in the northern part of the area, also give added irregularity to the otherwise uniform surface of the plain.

Character of the Drift

The drift in the Trenary till plain is fairly uniform in textural and other physical characters. Normally it possesses a fair association of clay, and constitutes a generally good agricultural soil. As a rule, the drift cover is thin, forming a thin veneer over the bed rock, and locally, especially in erosion channels, is lacking entirely.

In texture, the material of the drift is fairly heavy. The surface soil is loose and friable, and ranges from loamy sand to loam. The sub-soil is more firm and compact, usually quite heavy, and with a sandy clay character. Local patches of sand are distributed through the area but constitute a relatively small proportion of the surface. Ridges of sand and gravel, sorted, and in places well stratified, form a conspicuous feature in the relief. Drumlins, of the same material as the till plain, are found concentrated in the northern part.

The drift is commonly reddish in color and contains an abundance of large erratics. A large amount of locally derived limestone and dolomite

including material from the Hermansville, Black River and Trenton formations, is incorporated in the debris. Limestone is especially abundant in the southern part of the plain but decreases in amount northward, where the till becomes more silicious on approaching the belt of Cambrian sandstone. Near the Cambrian-Ozarkian contact, and for some distance southward, the till contains slabs of silicious dolomite carrying large amounts of the green mineral glauconite, described by Bergquist (10).

Much red sandstone, from the Lake Superior formation of Cambrian age, is intermixed with the drift in the northern part of the plain. It is probable that the red color in the soil throughout the plain is due to the incorporation of comminuted fragments of this highly ferruginous sandstone which was transported by the ice sheet in its southward advance.

The Eskers on the Trenary Till Plain

The numerous eskers, (Plate IV, Fig. 2) scattered over the surface of the plain, are composed of material ranging in texture from cobbles and coarse gravels to finer sands, rudely sorted and somewhat stratified. Blocks and slabs of local limestone are distributed through some of the ridges and occasionally form the bulk of the structure.

From the coarse nature of the included sediment it appears that many of the eskers, especially the shorter ones, were built up under conditions of rapidly moving waters in sub-glacial tunnels according to the theory as explained by Davis (11). The longer eskers were undoubtedly built up as a series of kames, by melt waters escaping into reentrants of the ice front, and became drawn out into long lines by the slow retreat of the edge of the ice while kame deposition was in progress, in accordance with the ideas advanced by Trowbridge (12).

The wide distribution of the eskers and their apparent lack of uniformity in trend and symmetry, together with the imbricated patterns developed, suggests stagnation of the Green Bay ice lobe to the north of the inner moraine, with consequent surface dessication. It is evident from the undisturbed condition of the stratified material contained in them, that the eskers were not formed in association with moving glaciers, but rather, under the influence of inactive ice. The conditions under which the Trenary till plain and its associated eskers were formed probably differ very little from those which Russell (13) describes in the development of eskers in the region of the Malaspina glacier in Alaska. Anderson (14) and Flint (15) have likewise emphasized the significance of eskers in the determination of stagnation areas in the waning stages of ice activity.

PLATE V



Figure 1.—Drumlin in Till Plain, one-half mile west of Eben, Alger County.



Figure 2.—Cross section of Drumlin showing heterogeneous arrangement of boulders and cobble in till.

Drumlins

A few drumlins are situated in the Trenary till plain in the vicinity of Eben Junction and Chatham (Plate V, Fig. 1). They are somewhat steep sided and elongated, with a roughly north-south trend, parallel to the direction of ice movement. Compared with drumlins in Menominee County, the drumlins in this area are poorly developed. They average about twenty to forty feet in height and from less than a half to more than a mile in length. The few drumlins in this district are composed of a reddish colored, loose textured, boulderly till and contain an abundance of locally derived, flat blocks of limestone and dolomite. There is no definite arrangement of the material in these ridges, the coarser erratics and limestone slabs being set into the loose textured matrix in all positions and orientations. (Plate V, Fig. 2)

The presence of the drumlins in the northern part of the till plain demonstrates a renewed activity and readvance of the ice for a short distance over the previously deposited till plain, according to ideas advanced by Russell (16).

Conclusions

The conclusions which are drawn in regard to the origin of the Trenary till plain may be summed up as follows:

(1) The plain was formed in the axis of the Green Bay lobe following its final recessional halt as marked by the position of the innermost moraine in this area.

(2) The till comprising the plain, was deposited largely under an ice sheet which became stagnant and melted en masse through surface ablation.

(3) The eskers distributed over the surface of the area were formed largely in tunnels beneath the stagnant ice mass, also, in part, through the recession of kames built up in reentrants to the lee of the glacier, by melt waters issuing from the ice during its marginal retreat.

(4) After the stagnant ice mass had melted down there was a slight readvance of ice, probably from the Superior lobe which, in the northern part of the plain, modeled the surface into a series of poorly defined drumlin ridges.

The Cooks Moraine

Immediately west of Indian Lake, in Ts. 41 and 42 N., R. 17 W., Schoolcraft County, stands an isolated remnant of a moraine which is probably associated with the Green Bay lobe. It appears to represent the front of an ice sheet which moved in a southeasterly direction and is related to the same movement which affected the Garden Peninsula, farther south,

in Delta County. The reason for this interpretation is to be found in the bearing of the striae which is S. 15 E. near Cook's Mill.

The bulk of the moraine stands at an elevation between 750 and 800 feet and, except for the marginal borders, was above the level of the Algonquin waters (770 feet at Cook's Mill). The drift in this area is relatively thin and consists of a clayey to loamy till, with numerous limestone blocks and erratics. The glacial cover ranges from 0 to 25 feet in thickness and rests directly upon a floor of limestone. The topography varies from fairly smooth in the southern part of the area to undulating and irregular in the north. In much of the region, especially where the till is thin, the surface expression of the terrane is a direct reflection of bed rock control.

If the correlation of this morainic remnant is correct, then it is obvious that it represents the oldest and outermost position of the Green Bay lobe in the Lowland province of the Northern Peninsula. Consequently, it was deposited at an earlier date than the oldest outer moraine which is associated with the Superior lobe.

Outwash Plain Associated with the Cooks Moraine

A narrow belt of outwash fringes the southeast border of the moraine in the vicinity of Indian Lake, Schoolcraft County. The feature is relatively smooth and breaks gently into the level sandy lake plain on its outer margin. Near the edge of the moraine the plain stands at an altitude of slightly more than 750 feet and was undoubtedly washed by the waves of the Algonquin water as they beat against the shore cliff of the plateau. The highest part of the outwash plain rises to the level of the water plane of Lake Algonquin (770 feet) and was consequently slightly inundated. The plain is composed of cobbly, gravelly and sandy material, shallow in depth and rests on bed rock.

FEATURES DEVELOPED BY THE SUPERIOR LOBE

Moraines

THE NEWBERRY OR "OUTER" MORAINIC SYSTEM

The final withdrawal of the ice sheet from the basin of Lake Michigan and its northward recession into the Northern Peninsula, marked the closing phase of activity of the Michigan lobe. No deposits which can definitely be related to this ice tongue are in evidence in the Tahquamenon-Manistique drainage area. Shortly following the disappearance of the Michigan lobe, and in the waning stages of the Green Bay tongue, the ice mass of the eastern part of the Superior basin readvanced over the area as an independent glacier. The early activity of this lobe, in the

PLATE VI



Figure 1.—A portion of the Steuben section of the Newberry Morainic System. Fire Tower hill, near Steuben. T. 44 N., R. 17 W.



Figure 2.—A closer view showing undulating character of topography.

region east of the meridian of Munising, is associated with a well-defined but much broken morainic system, which for much of its course follows the southern edge of the Manistique-Tahquamenon swamp. It pertains to a lobe which covered the greater portion of the Manistique-Tahquamenon drainage basins in Schoolcraft and Luce counties. In view of the fact that this morainic system is so broken in trend, it will be necessary to consider it on the basis of its several units or segments.

The Steuben Segment—An Interlobate Feature

This feature has its western terminus in the east half of T. 44 N., R. 19 W. in Alger County. It is here characterized by low crested ridges and knolls which are separated by intervening areas of relatively flat, sandy outwash. In the extreme southeastern portion of the moraine, south of McComb Lake, the surface assumes a definite knob and basin type of topography. Numerous lakes are set within the basins between the knobs. Thence, it continues in a southeasterly direction and enters Schoolcraft County in T. 44 N., R. 18 W.

From the northeastern corner of Delta County, a well defined morainic limb extends southward along the east side of the Sturgeon River. This extension of the moraine into Delta County is unquestionably related to the Green Bay lobe and suggests an intermediate position of the ice front later than the Cook moraine, but before it receded to the Au Train-Whitefish lowland and previous to the shaping of the drumlins near Eben Junction by a readvance.

The position of the Steuben segment of the Newberry moraine, in the reentrant between the Green Bay and Superior lobes, definitely identifies its interlobate relationships, and accounts for the apparent strong relief of the feature. The eastern edge of the segment terminates in a sandy lake plain through which the Indian River has cut its channel. This river flows eastward through a lowland depression across the main mass on the moraine and has been responsible for much dissection on the morainal slopes. (Plate VI, Figs. 1 and 2)

This fragment of the outer moraine is generally sandy in character but carries local patches of clayey till and numerous erratics throughout. The topography, in the main, is moderately rolling to undulating. In relief, the crest of the moraine stands at an elevation between 750 and 900 feet. It rises above the plane of the Lake Algonquin waters and shows little or no effects of water modification except along the lower marginal borders which were washed by wave action.

The Hiawatha Segment

A second or middle fragment of the Newberry moraine finds expression as a low, water worked feature in T.s 42 and 43 N., R.s 15 and 16 W. It

PLATE VII



Figure 1.—Morainic topography in the vicinity of Blaney. A portion of the Newberry Morainic System shows characteristic level relief of water worked feature.



Figure 2.—Water worked morainic area of the Newberry System, in the Twin Lakes country. Sec. 10, T. 45 N., R. 10 W.

extends through the settlement of Hiawatha and is separated from its eastern and western components by broad stretches of intervening low-land swamp and lake plains. The soil in this section ranges in character from a gravelly to a sandy loam and contains little if any clayey material. Erratics and cobbles are strewn haphazardly over the surface and in some of the lower border tracts protrude through the thin overwash of sand, giving evidence of a much greater extension of the moraine in pre-Algonquin times.

The moraine near Hiawatha is relatively low crested and stands at an average elevation ranging between 730 and 740 feet. The surface character is subdued and level and portrays very definitely the effects of Lake Algonquin wave action. It stands 60 to 70 feet above the swamp southwest of it. The low lying margins of the moraine have been reduced to a level plain and the bordering outwash has been greatly modified by lacustrine activity. The demarcation between moraine and the overwash of Chamberlin (17), is not always distinct or definite due to the mergence of the two features at a more or less common level.

The Blaney Segment

A few miles east of Manistique, the main highway, US-2, ascends the Nipissing shore scarp to a sandy plain which stands at an elevation of 685 feet. This lake plain extends eastward for several miles beyond the scarp. Here it breaks off into a series of level disconnected morainal fragments which may be traced intermittently along the southeastern border of the Manistique swamp to the vicinity of Blaney. At this point, the ridge gains considerably in strength and assumes a fairly definite morainal character. (Plate VII, Fig. 1) It continues thus in a general northeasterly direction through McMillan and Newberry, to the southeastern corner of Luce County where it breaks off into a sandy lake plain.

This section of the moraine is confined essentially within the Manistique-Tahquamenon basin, north of the Niagara cuesta. It was molded by the ice sheet into a mass of rolling drift with fairly strong expression but possesses no definite ridge development. The outer edge of the moraine, for much of its extent, was built up to the level of the Niagaran escarpment and in places stands superimposed upon its crest with altitudes up to 1000 and more feet. The inner margin, on the other hand, borders the marshy lowland and descends to 720 feet in some of the lower areas.

The eastern end of the moraine, in the meridian of Blaney, is relatively wide, and averages twelve to fifteen miles across, from north to south, where it extends into Mackinac County. It narrows gradually to the east, and leaves the southeastern corner of Luce County in a series of dis-

membered fragments. The topography of the moraine is, for the most part, rough and irregular. It is characterized by a knob and basin type of development. The knobs are usually not very high and the basins are relatively shallow. The highest knobs, located immediately south of McMillan, rise to an altitude of over 1000 feet, and are well above the highest level of Lake Algonquin. The largest basin in the outer morainic system is occupied by Manistique Lake and covers an area of practically fifteen square miles in Luce and Mackinac counties. North Manistique Lake likewise fills one of the larger basins in the area.

The glacial drift which constitutes the moraine is largely sandy, with local admixtures of clayey material. The soil of the sandy areas is loose textured, porous and open, and allows good drainage. The heavier phases are more compact and firm and tend to retain a greater amount of water. The more important areas of clayey drift are found in the vicinity of the Manistique lakes and south of McMillan. Foreign boulders and limestone slabs of local origin are abundantly scattered through the glacial drift, especially in the more strongly developed inner portions of the moraine. In the weaker outer borders, the drift is sandy to gravelly in texture and erratics are generally quite scarce. These lower areas have been modified greatly by the waters of Lake Algonquin, which at one time covered most of the surface of the moraine. The true morainic character has here been masked by the accumulation of a thin veneer of fine sediment which was washed down from the higher inner slopes. Post-glacial streams have locally eroded their channels below the surface overwash to expose the underlying bouldery till.

An illustration of the variations which may occur is well displayed in the morainic area south of Newberry. The northern, or inner border of the moraine in this region is fairly strong in relief and carries a sandy, bouldery drift. Southward, down the back slope of the moraine, the topography becomes more gentle and subdued. Here the waters of the bordering glacial lake were active in modifying the older features. Surface boulders and erratics are very scarce, but a large amount of cobbly and gravelly material is enclosed in the drift. On the southern, or outer border, the relief is smoothed down to a low plain and the morainic character is almost completely lost. A thin outwash of several feet of fine gravel and sand seems to cover the original morainic topography, the till of which is exposed only where the surface mantle of sediment has been removed by erosion, especially in the valleys.

The drift of this morainic segment varies considerably in thickness, depending largely upon the disposition of the underlying bed rock surface. Where the rock floor is high and ridged, as along the crest of certain portions of the Niagaran cuesta, notably in the southeastern part of Luce County, the superimposed drift is very thin. The relief in such

areas is controlled directly by the underlying rock surface. In areas where the rock floor is deeply channeled, and particularly to the north of the escarpment, the drift has filled in the hollows and basins to a much greater depth. (Plate VII, Fig. 2).

Deep wells at the Asylum grounds in Newberry, on the inner margin of the moraine, enter rock at 320 feet. Wells which furnish the municipal supply of water for Newberry are in sand and gravel at a depth of 90 feet.

Two recesses occur in the northern border of the moraine: one between McMillan and Newberry, and the other, four to ten miles east of Newberry. In each of these areas there is a thick deposit of red clay of lacustrine origin. An artesian well recently drilled in the S. E. $\frac{1}{4}$ of the N. E. $\frac{1}{4}$ Sec. 8, T. 45 N., R. 10 W. penetrated 80 feet of red clay, above 50 feet of sand and 2 feet of gravel before entering bed rock. In the S. W. $\frac{1}{4}$ of the S. W. $\frac{1}{4}$ Sec. 13, T. 45 N., R. 8 W. artesian water was struck below 53 feet of red lake clay. Near Soo Junction in Sec. 35, T. 46 N. R. 8 W. a well penetrated 172 feet of red, gummy clay before reaching rock.

In conclusion, it may be stated that the Newberry morainic system represents the position of the ice front during the maximum extension of the independent Lake Superior lobe which readvanced shortly after the break up of the main Michigan ice tongue. The final withdrawal of the ice from the moraine marks the initial stage in the surface development of the extensive Manistique-Tahquamenon basins. The morainic system was originally laid down as a continuous feature and during the early history of the basin drainage played an important role in confining the waters of the lowlands behind it. The continuity of the morainic system was broken, first by the action of Lake Algonquin waters, and finally by erosion of through-flowing rivers which developed as uplift in the area progressed.

In terms of geologic time, the activities of the ice sheet which shaped the Newberry morainic system, may be correlated with the Sturgeon River moraine of the Green Bay lobe in Delta County. This would set the time, as following the complete disappearance of the Michigan lobe but antedating the formation of the eskers on the Trenary till plain. It seems probable that stagnation of the last remnant of the Green Bay lobe in the Northern Peninsula did not take place until the Lake Superior ice had withdrawn completely from the Newberry moraine and receded to a new front farther north.

THE MUNISING OR "INNER" MORAINIC SYSTEM.

A strong morainic system, formed by the Lake Superior lobe, and marking its second important halt in recession, may be traced from near

PLATE VIII



Figure 1.—High morainic bluff in the Munising System,
near the city of Munising.



Figure 2.—Moraines of the Newberry System, near Laketon,
S. E. $\frac{1}{4}$, Sec. 36, T. 46 N., R. 12 W.

the axis of the Green Bay lobe in the vicinity of Au Train Lake on the west side of Alger County, northeastward to the meridian of Grand Marais. Here it swings southeastward into the Tahquamenon swamp in the central portion of Luce County. For most of its distance in Alger County, the moraine parallels the shore of Lake Superior, although in places it lies some distance back.

The extreme western end of the system, where it swings southward into the Au Train-Whitefish depression, merges as an interlobate feature with the eastern limb of the inner moraine of the Green Bay lobe. In this section of the moraine the topography varies from rolling to gently undulating and in places flattens out to a fairly level surface. Erratics and boulders are prevalent and much sandstone and dolomite of local derivation are incorporated in the sandy gravelly clay till.

A narrow strip of moraine, not exceeding an eighth of a mile in width, borders the north edge of an extensive outwash apron from the west central portion of Sec. 31, T. 46 N., R. 19 W. for a distance of seven miles to the western edge of Sec. 19, T. 46 N., R. 18 W. It is separated from the main moraine to the north by a narrow bench of sandy lake plain which swings into the area from the southeast. This narrow belt of moraine appears to be an ice contact border and marks the position of the ice front during the interval when outwash material was being carried southward by melt waters. The morainal character of this contact belt is well preserved by the presence of the erratics and boulders incorporated, and by its relief.

The bench of sandy lake plain, which lies between the narrow contact belt and the main moraine to the north, evidently represents a fosse which was deepened sufficiently to allow the lake waters to remain after the main mass of the moraine had become uncovered.

In the vicinity of Munising the moraine shows considerable strength and is surmounted with high knobs which reach a maximum elevation of approximately 1010 feet. (Plate VIII, Fig. 1) Much of this area stood above the water of Lake Algonquin in its highest stage and was unaffected by its activities. The intercalation of lacustrine clay and sand with the glacial till on the morainal slopes is apparent to an altitude of 940 feet, or 340 feet above the datum plane of Lake Superior. Leverett (18) describes storm beaches on Scaffold Hill, in the vicinity of Wetmore, as high as 960 feet. He accounts for the water body being held up to this plane on the assumption that "the ice was still present in the great Manistique Swamp, to the east and southeast". Above 960 feet in this area, the slopes and crests are rugged. Below this level, however, the slopes are toned down and smoothed off considerably from the effects of water action along the Algonquin lake shores.

In the Munising-Wetmore portion of the moraine, the topography ranges from rolling and undulating in the higher positions, to gently wavy and somewhat flattened surfaces in the water-modified lower portions. The drift varies all the way from clayey till of ice deposition to local pockets of stratified sand and gravel of lacustrine origin in the modified areas. Erratics are abundant and, together with blocks of local sandstone and dolomite, constitute a considerable portion of the drift.

East of Munising, the moraine becomes quite bulky and reaches a maximum width of 14 miles in the meridian of Melstrand, where it breaks off into the marshy lowlands of the Cusino swamp. This section of the moraine is undulating in character and locally somewhat rugged in relief. It is quite generally creased by drainage channels with swampy borders which break up the continuity of the system on its eastern edge. The flattened crests and smooth surface slopes are significant of the activity of Lake Algonquin waters which completely covered the moraine.

From Melstrand, eastward for a distance of 17 miles, the moraine parallels the shore of Lake Superior as a tenuous belt, skirting the southern edge of the Beaver Lakes basin. The feature is extremely narrow for much of this distance, but is wider in places, ranging from less than one-fourth to several miles in width. The topography is irregular, beset with low hills on the inner border but flattens out at the outer or outwash margin. At the outwash border, the ridge has an average elevation of 800 feet but rises to almost 1000 feet in the more rugged portion directly south of Beaver Lake. Erratics and local boulders are numerous and the drift is sandy to loamy in character, with local admixtures of clayey till.

The general nature of this section of narrow moraine seems to indicate quite conclusively that it marks an ice-contact border. Here the margin of the glacier rested for a considerable period, during which it melted and spent its material in building up an outwash apron rather than a bulky moraine. That the ice front remained for some length of time in this position is evidenced by the vast extent of outwash gravel which was spread out south of the ridge. Leverett¹ confirms the writer's view that the ice must have formed a contact border in this area.

In the meridian of Au Sable Point in Alger County the Munising morainic system is offset to the south about three or four miles. Here it sets in again as a bulky mass of rolling drift and attains a maximum width of 10 miles where it extends southward into Schoolcraft County in R. 14 W. Near Grand Marais the inner border is almost two miles south of the lake shore. In this vicinity its continuity is broken by a low lying, swampy, outwash re-entrant, which extends into the Sucker Lake

¹Leverett, F.,—Personal communication.

area in Schoolcraft County. This strip of poorly drained outwash may be traced northward into Alger County for a distance of seven miles through T. 40 N., R.s 13 and 14 W. Highway M-77 follows this lowland between the morainic segments for some distance, in the region from Seney to Grand Marais.

In the eastern end of Alger County the morainic segment exhibits an undulating to rolling relief. It ranges in altitude from 800 feet, along the borders of the outwash plain on the south, to over 1000 feet in the southwestern corner of Alger County. Much of its surface particularly below 950 feet was inundated by the waters of Lake Algonquin and presents definitely the results of wave activity in washed down slopes and subdued crests.

The drift in this area is composed predominantly of a grayish colored loamy sand, intercalated with lenses of reddish-gray silt and fine sand. Boulders of local sandstone and erratics are embedded in the loose till matrix.

In the region immediately west of Beaver, situated on M-77 about ten miles south of Grand Marais, the moraine loses strength. It was here modified to a considerable degree by the waters of Lake Algonquin. The surface has been eroded and reduced almost to the level of a flat topped plain. Small, scattered hills of low relief, together with erratics and the presence of numerous kettle basins, give support to its morainal origin. Along the Adams trail, west of Beaver, the outer margin of the moraine has been so completely subdued by lacustrine erosion and subsequently mantled with thin lake deposits of sand, that it is difficult indeed to discern its till character. However, in places, the till with its associated boulders, projects through the thin veneer of lacustrine sand and makes possible the true interpretation of the feature. The morainic expression becomes more pronounced towards the northern inner border.

Upon entering Luce County, the Munising morainic system takes a definite swing to the southeast and breaks off into the Tahquamenon swamp a few miles north of Dollarville. Like its counterpart south of the swamp, this moraine is not separable into distinct ridges, but stands as a mass of undulating drift.

On its inner border, the moraine here stands at an altitude of 750 to 800 feet and presents a strongly developed relief. The topography is generally quite undulating and in places broken by numerous basins which are hemmed in by knobs of varying heights. The drift is predominantly sandy and is locally interspersed with admixtures of gravel and clay. Erratics and occasional slabs of limestone are scattered through the section but not in such large quantity as in the moraine to the south. The limestone makes up a relatively small proportion of the drift but is conspicuous in the shore bluff which marks the north-

ern margin of the moraine, south of Cold Spring in Sec. 6, T. 47 N., R. 10 W. This shore feature marks the northern limit of limestone drift in Luce County and is undoubtedly quite near the Cambrian-Ozarkian contact.

The outer border of the moraine is relatively low and exceedingly weak. It descends into the Tahquamenon swamp at altitudes ranging from 700 to 720 feet. The drift here is largely sandy with gravel intermixtures and with but few erratics exposed. The surface is smooth and unbroken, except where it has been seamed with drainage creases which have cut down into the unmodified drift below.

This section of the morainic system was largely inundated by the waters of Lake Algonquin. A small area covering approximately ten square miles, in the southeastern part of T. 48 N., R. 12 W. and the northwestern corner of T. 47 N., R. 12 W. in Luce County, extends westward into the northeast corner of Schoolcraft and the southeast corner of Alger counties, where it rises to an altitude of 1000 feet. Here it was above the Algonquin level and stood as an island mass in the lake during the time of the greatest expansion of the lake. Six miles of marshy lowlands separate the eastern border of this segment of the morainic system from its eastern continuation in Luce County. This fragment of the moraine is completely enclosed by marshy lowland but may be traced as a conspicuous feature for a distance of eight miles to the depression of the Tahquamenon river, in T. 48 N., R. 8 W. East of the river the morainic system leaves the County in the vicinity of the Big Falls and enters Chippewa County twelve miles south of the Lake Superior shore. In Chippewa County it continues eastward, skirts the south shore of Whitefish Bay and finally merges with the Newberry moraine near Eckerman. From this point the combined systems follow through to Nadoway Point, the easternmost terminus of the moraine.

Significance of the Munising Morainic System

Although the present morainic system is broken into a number of units widely separated by rivers and marshes, it is nevertheless, obvious that it may have extended across the region with a continuous front shortly after the ice sheet had retired. It marks very definitely the border of the Superior lobe at the time when the Green Bay tongue was holding its final position in the peninsula. It is quite probable, that at this time the Green Bay ice sheet made the readvance that shaped the drumlins near Eben Junction and Chatham.

The position of the ice, as it stood upon the Munising moraine, was such as to block the present outlet of the Tahquamenon river in northeastern Luce County. Border drainage from the melting ice front was carried southward across a low plain which apparently had received

a very small contribution of drift from the retreating glacier. The discharge waters, flowing naturally toward the Lake Michigan basin, were for a time, undoubtedly impounded behind the as yet unbroken outer Newberry morainic system. Thus a more or less extensive interior lake developed between the outer and inner moraines, and persisted for some little time until the water succeeded in cutting channels across the barrier. At this stage, a number of rivers, extending from the outer margin of the Munising moraine and flowing into this interior lake, were formed upon the initial, slowly uplifting slopes of the intermorainic plain. The Tahquamenon drainage basin was as yet undifferentiated but in all probability received its due proportion of melt-water drainage from the waning ice.

THE CRISP POINT MORAINE

The final position of the Lake Superior lobe and of the Wisconsin ice sheet as a whole in Michigan, is marked by a narrow, transected morainal ridge of relatively weak expression, trending across the north edge of Luce County a few miles south of the Superior shore. It crosses the county in a series of broken fragments, which in some areas have been so completely dissected by wave erosion as to make it difficult to trace.

The westernmost fragment of this morainic ridge lies south and west of the Sucker River basin, in the western part of T. 49 N, R. 12 W., where it extends for a few miles west into Alger County. In this section the moraine has a fairly rugged development but does not stand sufficiently high to have extended above the waters of Lake Algonquin. Hence, it presents a water-washed and much modified surface. The expression is typically of the knob and basin type with many of the basins without lakes or swamps.

A second fragment continues eastward from the eastern end of the Sucker River basin through T. 49 N., R. 11 W. and thence into R. 10 W. The inner border of this section is quite definitely marked by a conspicuous, although somewhat broken, wave-cut shore which follows along at 715 feet elevation. The crest of the moraine, slightly south of the shore cliff, rises to an altitude of 730 to 750 feet but possesses a decidedly subdued relief. In the northwestern portion, however, its surface is deeply basined and the topography is rugged, due to the melting in place of detached blocks of ice which were left in the wake of the retreating glacier. The outer or southern border, on the other hand, is not so distinctly marked. It merges more or less indefinitely with the adjoining lake plain and becomes difficult to differentiate. The drift in this morainal segment is sandy, light, and loose textured. It contains scattered patches of gravel, cobble and occasional small amounts of clay. Boulders are relatively scarce although locally they may be quite evident.

In the northeastern corner of Luce County, extending east of the Two Hearted River into Chippewa County, near Crisp Point, is another fragment of the ridge. Here the moraine has developed a fairly strong expression and has a relief much higher than its western counterparts. The crest rises to an altitude of 800 feet but still stood below the highest Algonquin water plane. The surface is undulating and characterized by a swell and sag topography. The drift is sandy to clayey in texture and contains numerous boulders of fairly large dimension. The north edge of this morainal remnant is marked by a wave-cut shore cliff standing at a general elevation of 715 feet. Southward, the moraine loses its ruggedness and on its outer border merges indefinitely into the sandy lake plain and swamp of the Betsy Lake depression.

The general trend of the moraine, as evidenced by its various fragments from west to east across Luce County, suggests a north-northwest recession of the Wisconsin ice border from the southeastern end of the basin of Lake Superior. It is the youngest glacial feature in Michigan and marks the last definite position of the continental glacier in its retreat from the state. It is obvious from the weak expression of the moraine, that the ice which formed it did not hold its front in position for a very long period before finally melting.

The stagnant remains of the Green Bay lobe in the Trenary till plain region in Alger County, were no doubt, dissipated before the Superior ice sheet had reached the position of the Crisp Point ridge.

Outwash Plains

During the process of moraine construction, the ice borders remained somewhat stationary for long intervals, due to the established balance between advance and retreat. In the dissipation of the glacier front, a large amount of melt water was carried away much of the time, but more especially in the summer when melting was rapid and on a larger scale. Thus, water work accompanied ice work in all cases of melting and resulted in the modification of much of the material previously deposited by the glacier.

The glacier streams, which flowed from the melting ice across the moraines, carried considerable quantities of sediment out to positions in front of the borders. In the areas where the outflowing streams emerged upon more gentle slopes, velocities were reduced and the material carried was deposited in the form of alluvial fans. Fans from adjoining stream courses, building out in all directions, in places merged with each other to form broad alluvial plains of stratified gravel and sand. These plains were generally constructed in front of the moraines in the form of outwash and overwash aprons.

Outwash plains or aprons are, as a rule, composed of both coarse and fine material. The outflowing streams of melt water, transporting debris from the ice border were forced to drop the gravel and cobbles almost immediately upon reduction of velocity. The finer material, however, was distributed for considerable distances from the source of the sediment. Thus, the load carried was sorted out according to texture and rearranged by the water into stratified deposits, with the coarsest material spread out nearest the retreating ice front.

Features of this type are usually characterized by gentle to moderate relief, except where pits have been formed by the melting of detached ice blocks below the surface. The topography may then assume an irregular aspect but maintains, nevertheless, a generally even skyline. The material of outwash plains ranges from coarse silt to sand, gravel and cobble of varying textures but usually with a preponderance of the less coarse assortment. The structure is loose and porous, with the result that drainage is excellent. Erratic boulders may be present near the morainic margins of outwash plains, but are inconspicuous in other portions save where the mantle is so thin as to allow the shallow underlying till to protrude.

Outwash plains constitute a relatively small portion of the surface area in the Tahquamenon-Manistique drainage basins. They are confined largely to the outer borders of the Munising moraine and occur more or less as widely separated or isolated features. Like the moraines, the lower portions of the original outwash plain have, in part at least, been greatly modified by the activity of Lake Algonquin waters and, in certain areas, the feature has been completely removed by prolonged wave erosion.

OUTWASH PLAINS ASSOCIATED WITH THE MUNISING MORaine

The Wetmore Plain

A conspicuous outwash apron extending from the Au Train-Whitefish depression in Alger County, eastward for a distance of ten to twelve miles into Schoolcraft County, occupies a position in the reentrant between the Green Bay and Superior lobes. It was built up in the course of ice recession, to a general level of close to 1000 feet near the ice contact, and except for its outer marginal borders stands well above the plane of Lake Algonquin. It is composed of loosely consolidated and porous gravel and sand. The surface slopes gently to the south-southeast, a condition largely controlled by its interlobate position. It is pitted with numerous basins, many of which hold shallow, irregular-shaped lakes. These were undoubtedly formed where detached masses

PLATE IX



Figure 1.—Burned over outwash plain. south of Kingston Lake.
T. 48 N., R. 15 W.




Figure 2.—Section of outwash plain showing stratification in the sandy
material (Mattock 14" long). Sec. 4, T. 47 N., R. 13 W. Along M-77.

and blocks of ice remained after the active ice border had retreated beyond them.

The Wetmore plain has an average width of 5 to 7 miles through much of its course. It is flanked on its inner margin by a narrow fringe of moraine which appears to mark the position of an ice-contact border. The pitted nature of the surface, together with the unusual width of the plain, in contrast to the narrow associated moraine, seems to support Leverett's (19) views that the feature was built up in the course of ice recession.

From the meridian of Wetmore, eastward for a distance of twelve miles, to the vicinity of Melstrand there is little or no outwash fill along the outer border of the Munising moraine. Leverett (20) attributes the absence of outwash in this area to the ponded condition along that part of the ice border. He states also that "the parts where outwash plains occur were high enough to stand above Lake Algonquin or were very close to the upper level of the Algonquin waters." It seems probable, however, that the absence of outwash fill in this area may be accounted for by simple removal by wave action of material not built up sufficiently high to withstand the ravages of lacustrine erosion.

The Boot Lake Plain (An Outlier of the Wetmore Plain)

About five miles south of Shingleton, T. 45 N., R. 17 W., in Schoolcraft County, the main highway, M-94, rises to a gravelly, cobbly table land which stands at an altitude of 820 to 850 feet. This isolated patch of outwash, bordered on all sides by marshy lowland, covers an area of practically 10 square miles. It breaks abruptly, with steep escarpment slopes on all sides, to the surrounding marginal swamp. The plain is confined largely within the inter-stream area of two headwater branches of Stutts Creek, which apparently were responsible for its isolation and the development of its marginal cut-cliff character.

This small area of outwash plain is extensively pitted and carries a number of shallow, irregular basined lakes among which, Lilley, Boot, Sand, and Clear lakes are the most important. The elevation of the surface of this plain is lower than the east edge of the Wetmore plain, which is about six miles west, but lines up with it quite definitely when surface slopes are considered. The general altitude of this outwash fragment suggests that it is an outlier of the Wetmore Plain which was first isolated in part by wave action of Lake Algonquin waters and later by the erosional work of the post-glacial streams which now almost completely surround it.

This table-land, like its main western counterpart, stood slightly above the plane of Lake Algonquin and was not subjected to the modifying activity of the lake waters.

The Kingston Plain

A rather extensive outwash apron may be traced along the north edge of the Manistique-Tahquamenon swamp from the northwestern corner of T. 47 N., R. 16 W., in Schoolcraft to the west edge of Section 12, T. 46 N., R. 11 W., in Luce County. (Plate IX, Fig. 1) This plain covers an area of several hundred square miles and is by far the largest single outwash feature in the region. A tongue of the plain in Alger County, extends northward into the reentrant between two morainic segments in T. 48 N., R.s 13 and 14 W. M-77, the main trunk highway between Seney and Grand Marais, crosses this portion of the plain for a distance of about 8 miles. In this area, the plain is considerably broken by low-lying, poorly drained, swampy depressions, and is represented essentially as sandy, tree-covered ridges separating the undrained lowlands.

In Alger County, the plain is bordered on the north by a narrow belt of terminal moraine which undoubtedly represents another ice-contact border. It does not seem justifiable that such an extensive outwash apron as has here been developed could have formed from a moraine so narrow, unless the ice front remained stationary for long periods and during the continued wasting of its front, discharged much of its debris into the outflowing streams of melt water. Thus, when the ice sheet finally melted in retreat, there was very little debris remaining in the mass to be dumped, with the consequent result that a very narrow belt of moraine was left to mark the contact position of the ice.

With the possible exception of the area in Alger County immediately bordering Kingston Lake on the east and south for a distance of several miles from its margin, the plain is uniformly level and presents a relief which is characteristic of such features. The relatively even surface is etched in places with shallow lakes and depressions which give a pitted expression to the topography. These pits suggest that much of the outwash was formed in the path of the retreating ice margin.

East and south of Kingston Lake, however, the surface expression is irregular and broken, and gives the impression of a morainic relief. The topographic irregularity is due to deep pockets and basins which have been extended downward rather than to elevations developed above the general plain level. The sky-line is even and smooth over the area, despite the roughened surface character. This portion of the plain may owe its development to deposition of gravels and sand over a nearly continuous sheet of stagnant ice of irregular thickness, as suggested by Thwaites (21). Or it may have assumed its surface character through the melting of large detached blocks of buried ice which resulted in slumping of the surface.

Thwaites (22) regards the presence of pitted outwash in front of a moraine as "indubitable proof that the glacier extended beyond the

moraine at a date not many years before its formation." Likewise, "a moraine behind a well developed pitted plain is the product of a comparatively rapid recession of the ice front and not of a readvance of any considerable magnitude." The conditions as found in this locality seem to justify these conclusions. The lack of erratics and glacial boulders in this area seem further to bear out the opinion that the feature is outwash rather than moraine. It also manifests a sufficiently thick overwash deposit of glacio-fluvial material to have completely buried any remnants of a moraine which may previously have been deposited.

The material of the outwash varies somewhat in the different areas. In the inner margin, near the morainal contact, the sandy-gravelly drift is intermixed with cobbles and coarser debris which would indicate transportation by rapidly flowing melt-water streams. Southward, extending into Schoolcraft County, the deposit assumes a more uniform texture ranging from sand to gravel, and is stratified in character. (Plate IX, Fig. 2)

Lakes and swamps are numerous and widely distributed throughout the extent of the plain, but on the whole the drainage is easy and follows southward down the natural surface slopes.

The eastern extension of the plain, continuing into Luce County, and especially that portion situated directly across the swamp from the village of McMillan, seems to have formed in the wake of a rapidly retreating ice-border. Near its inner edge, the plain is generally smooth and unbroken. It contains coarse cobbly and gravelly material interspersed with occasional boulders which protrude through from the underlying till. To the south, however, the surface of the apron is deeply trenched and pitted and presents a relief sufficiently rugged to be easily mistaken for a morainic feature. Superficial boulders are relatively rare in this area but may be encountered within a few feet of the surface as evidenced in the deeper skidding trails and along the floors of streams which have trenched their channels through the overwash.

It appears that this apron was formed upon a more rugged morainic topography, during a recession of the ice border which was marked by a succession of short halts. The outflowing melt-water streams did not carry out a sufficient amount of sediment to completely fill the depressions and smooth out the irregularities of the original surface. Leverett (23) suggests that the plain was here built up in a series of steps from south to north as the ice retreated from the area. He states further that "the ice contact at the north edge of each plain is marked by a low bluff-like rise, the northern edge of each plain being a little higher than the southern edge of the next one and being trenched in places by the passage of streams across it from the next later plain."

Chapter IV

LACUSTRINE GEOLOGY

GENERAL DISCUSSION

For a considerable time after the final disappearance of the Superior lobe from the position of Crisp Point Moraine in Luce County, and while the Wisconsin ice sheet still blocked the North Bay outlet in the northeast portion of the Superior Basin, there continued a widespread inundation of the Lowland province by the waters of Lake Algonquin. The greater portion of the peninsula from the meridian of Munising eastward was flooded, and remained so until the retreat of the ice fully uncovered the Ottawa drainage line. Only a few small isolated areas in this region rose above the level of the highest Algonquin waters and they were represented merely as islands.

Leverett (24) states that "in the southern part of the Peninsula near Gladstone, and on the Garden Peninsula east of Big Bay de Noc, the altitude of Lake Algonquin was about 120 feet above Lake Superior, but at Marquette it was 340 feet and just north of Sault Ste. Marie it was fully 400 feet, while at the southern end of Lake Huron at the head of the St. Clair outlet the altitude was only five feet above Lake Superior."

The glacial lake history of the Northern Peninsula is associated primarily with developments which took place in the Lake Superior basin and to a less extent with those in the Lake Michigan depression. According to Leverett (25), "the oldest and highest of the large glacial lakes in the Northern Peninsula was held in the basin of the Ontonagon River to the south of Copper Range. During the existence of this lake the southern edge of the ice sheet appears to have rested on the Copper Range so that the waves were beating against an ice barrier on the north side of the lake." He places the altitude of this initial lake, Lake Ontonagon, at 1320 feet above sea level or more than 700 feet above Lake Superior and indicates that, during its short existence, the line of discharge led west from Lake Gogebic.

Lake Duluth, which followed Ontonagon, began as a small body of water in the extreme western end of the Superior basin, when the ice in that area had melted back for a short distance. This lake drained southward through the Brule River into the St. Croix and thence into the Mississippi. Leverett places the outlet through the Brule at 500 feet above Lake Superior in the beginning stages of the lake. During

its history, the outlet was cut down 50 feet and the water plane was lowered correspondingly.

The waters of Lakes Ontonagon and Duluth were confined, by the position of the ice barrier, entirely to the highland area in the western part of the peninsula. They had no direct influence upon developments within the Tahquamenon-Manistique basins inasmuch as the main mass of the Wisconsin ice sheet covered this area at that time.

As the ice in the Superior depression melted back eastward, a larger portion of the basin was uncovered. The lake waters expanded and at the same time the level was dropped to a plane below the Brule outlet. Thus there gradually came into existence a lake which eventually became confluent with the waters of the Huron and Michigan basins in the establishment of Lake Algonquin.

LAKE ALGONQUIN

Because of its widespread effects and influences upon the Tahquamenon-Manistique area, it will be necessary here to consider the problem of Lake Algonquin in some detail. In many respects, however, the details can never be complete because of the fragmentary evidence at hand in many of the areas. Shore line and strand features may be traced somewhat interruptedly for many miles and then again they may be entirely missing for equally long distances, with gaps between merely to be interpolated.

Field evidence available in Alger and Luce counties seem to indicate that the complete development of Lake Algonquin from the Duluth stage did not occur in one single change. Rather, the transition was gradual and during the process there were various incidents, as may be judged from the changes in the drainage lines of discharge.

One of the earliest connections of drainage between the Superior and Michigan basins, as correlated with Lake Algonquin, is associated with the Au Train-Whitefish depression, the head of which is situated in Au Train Lake, 9 miles west of Munising. During Algonquin times, when the border of the Wisconsin ice was standing in the vicinity of Munising and Grand Island, the water plane was raised to the level where it found an outlet through this passage into Little Bay de Noc. Although this channel has not been studied in complete detail in all of its course to Lake Michigan, sufficient evidence is at hand in the Alger County section to draw conclusions concerning its relations to drainage. Winchell (26) makes brief mention of a portion of this outlet in one of his early papers.

During the interval of discharge through this channel a large amount of water must have been carried with great velocity. The floors of both the Au Train and Whitefish rivers show evidence of considerable scour-

PLATE X



Figure 1—Algonquin terrace and wave-cut cliff, one mile South of Chatham



Figure 2.—Upper Falls on the Au Train River, near Forest Lake.

ing action. Bed rock, including the Hermansville dolomite in the north and Trenton limestone in the south, is exposed in extensive portions of the channels. Wherever drift occurs, it is generally so shallow that its base may be easily reached with a four-foot soil auger.

The drainage through this outlet was for a period sufficiently long to allow the formation of well defined wave-cut shore cliffs. These stand, in the region directly east of Limestone, at an altitude of 850 feet and may be traced northward out of the channel into the wide funnel opening a mile south of the village of Chatham. (Plate X, Fig. 1)

The present drainage basin carries two rivers: the Au Train flowing north into Lake Superior and the Whitefish which flows south into Lake Michigan, entering at Little Bay de Noc, near Rapid River. The divide between these two rivers stands a short distance south of Mud Lake in Section 30, T. 45 N., R. 20 W., at an altitude of 768 feet above sea level. Here the East Branch of Whitefish River gathers its head-waters for the Whitefish River farther south. The Au Train River draws its head-water supply from Mud Lake and a large number of feeder springs. In its northward course, near the village of Forest Lake, the river plunges almost 100 feet over a series of falls and cascades which have been carved out of the Hermansville formation, described by Bergquist (27). (Plate X, Fig. 2) It enters Au Train Lake at the Nipissing level and flows out across its sandy lake plain in a series of broad meanders, induced by low gradient, into Lake Superior.

In discussing the Algonquin beach in the Au Train-Whitefish lowland, Leverett (28) states as follows: "From a point near the Delta-Alger county line northward to Chatham there is a rise of 93 feet in a distance of 13.5 miles, or about 7 feet to the mile. In the southern 8 miles of this line the rise is 65.5 feet or slightly more than 8 feet to the mile, and in the northern 5.5 miles the rise is 27.5 feet, or 5 feet to the mile. This rapid rate of tilting is in a narrow passage only 1 to 1½ miles wide. Farther north, where the rate is lower, the waters had greater width."

Another passage which was opened up between the Superior and Michigan basins extends from Munising, southeastward through Anna River into the head of the Manistique swamp. This outlet was established at a time when the ice in the Superior basin had receded eastward beyond the meridian of Wetmore. The altitude of the passage is 838 feet, as determined by Leverett¹ by means of levels. The crest of this drainage way stands about 70 feet higher than the divide in the Au Train-Whitefish lowland. Hence, it probably functioned as an outlet for a short period during the early stages of outflow in the former channel, at a time when the ice barrier held the water plane to the higher level. The Anna River valley has been cut deeply through the Munising

¹Leverett, Frank, Unpublished field notes.

morainic massif which in this region exceeds 1000 feet in altitude. The depression is narrow, however, and shows little evidence of excessive scour. It is obvious from these facts that the waterway functioned as a connecting passage for a comparatively short period.

Lacustrine activity, in the region of Munising and Wetmore, may be traced to an altitude of 950 feet. The slopes of the high moraines in this district show the effects of water action almost to the crests where lacustrine silts and clays are intercalated with the glacial till. Leverett (29), reports gravelly bars at 946 and 950 feet about a mile north of Wetmore and a storm beach at 960 feet on the north slope of Scaffold Hill south of the station.

These beaches are much too high to fit the highest Algonquin plane which should here have an altitude of 875 and 880 feet. It is probable then, as suggested by Leverett, that the higher beaches in the region are the product of local waters rather than the waters of Lake Algonquin. The high level of the water during this stage may be accounted for on the hypothesis that a large mass of stagnant ice still persisted in the Manistique swamp to the east and southeast and held the water up.

Still later, after the ice had retired from the land surface of the peninsula and the greater portion of the Superior basin was uncovered, a new passage was opened between the Superior and Michigan basins by way of the Tahquamenon-Manistique drainage depression. At this stage the glacier front blocked the North Bay outlet in Ontario. The drainage was affected through a narrow strait extending from the vicinity of Dollarville westward to $1\frac{1}{2}$ miles beyond Danaher. At Dollarville the passage is about 3 miles wide and is bordered on both north and south by high moraines. Four miles westward, the channel widens out to 7 miles and includes a broad reentrant in the inner border of the Newberry moraine where a considerable formation of red lacustrine clay has been deposited. At McMillan the passage is $3\frac{1}{2}$ miles wide, but narrows again to less than 2 miles in its narrowest portion, near Danaher. For the entire distance of 13 miles between Dollarville and Danaher the narrow channel is bordered on the south side by the massive Newberry morainic system whose bulk and altitude prevented undue widening in this area.

A mile and a half west of Danaher the passage opens up into the broad Manistique swamp which spreads out over the greater portion of the surface of Schoolcraft County. The headwaters of the Tahquamenon and the East Branch of Fox River, a tributary of the south-flowing Manistique, extend into the area and are separated by the low divide which stands in the swamp immediately north of Danaher in Section 27, T. 46 N., R. 12 W. The altitude at the divide is 720 feet above sea level

and represents the lowest possible connection between the Lake Superior basin and the Manistique drainage area.

The opening of the Trent outlet at Kirkfield, Ontario, caused the level of Lake Algonquin to become fixed. Taylor places the upper strand of the beach at Kirkfield at an altitude of 883 feet above sea level, or 276 feet above the Algonquin beach in the area of horizontality. During the Kirkfield stage of Lake Algonquin the entire area of the Tahquamenon-Manistique drainage ways was inundated, and it was during this interval that the various morainic systems of the Lowlands province were modified so effectively by wave action.

The Trent Valley functioned as an outlet for Lake Algonquin until the area was uplifted to such a height that the lake could no longer discharge through it. The overflow was then diverted to the Port Huron and Chicago outlets. In the transition from the Kirkfield to the Port Huron-Chicago stages of drainage development, there was an interval of short duration when the three-outlet phase was active, but as the uplifts continued to the north, the Kirkfield outlet was finally abandoned. The waters discharged through the Port Huron outlet were carried into the Erie basin and at Chicago they were drained through the Desplaines, Illinois, and Mississippi rivers to the Gulf of Mexico, as described by Alden (30). According to Taylor (31), "it was during this stage that the main part of the great northern uplift took place, raising the region north of Lake Huron, and Georgian Bay and most of that around Lake Superior by at least 600 or 700 feet."

It was at the time of this two-outlet phase that the water level of Lake Algonquin began to fall. Uplift movements to the north and the opening of lower outlets to the south and east resulted in a definite lowering of the water plane and the development of the lower Algonquin beaches commonly referred to by Leverett and Taylor as the Battlefield and Fort Brady shores.¹ The flood waters in the Tahquamenon-Manistique drainage basin probably separated in the vicinity of the divide near Danaher prior to the latest Algonquin (Fort Brady) episode. Part of the drainage was directed northward into the Superior basin and the remainder was diverted southward in the direction of the Michigan basin. The McMillan straits then ceased to function as a waterway connection.

¹In U. S. G. S. Monograph 53—The Pleistocene of Indiana and Michigan, Leverett and Taylor apply the name Battlefield (pp. 435-436) to that well defined shore on Mackinac Island which stands at an altitude of 715 feet, and the name Fort Brady (pp. 436-38) to the series of lowest Algonquin beaches which stand at altitudes of 639, 653, 661 and 681 respectively.

From his studies of shore lines on Isle Royale and the Georgian Bay area, George Stanley is disposed to make the Battlefield and Fort Brady beaches in the vicinity of Grand Marais higher than mapped by Leverett and Taylor. He has found that the Fort Brady beach on Isle Royale rises rapidly to the eastward and so should be made much higher than previously indicated. At the time of this writing (August 1936) the issue as to the exact correlation of these beaches is still controversial.

PLATE XI



Figure 1.—Algonquin shore ridge, at 870 feet A. T., against the remnant of morainic ridge, Luce County. N. W. $\frac{1}{4}$, Sec. 10, T. 45 N., R. 10 W.



Figure 2.—Closer view of shore shown in Figure 1.

Further temporary halts in recession of the waters in Lake Algonquin are marked by a succession of beaches which in the region to the south of Grand Marais, in Alger County, may be correlated with the lower Algonquin stages. (Battlefield and Fort Brady)

When the ice sheet finally withdrew from the Mattawa and Ottawa valleys in Ontario, the North Bay outlet was opened. Thus, much of the overflow from Port Huron and Chicago was diverted to the north, with the result that the outlet was excavated to the rock sill at the North Bay col. The lowering of the outlet at the divide, together with the complete withdrawal of the ice from the basin, marked the close of the glacial Lake Algonquin and the inauguration of the non-glacial Lake Nipissing.

Shore Features Pertaining to Lake Algonquin

HIGHEST ALGONQUIN SHORE IN THE TAHQUAMENON REGION

The highest shore work recorded in the Tahquamenon area of Luce County is to be found in the high moraine immediately south of McMillan. The slopes of the moraine in this area show the effects of wave washing up to an altitude of 870 feet. This places the highest Algonquin level about 30 feet below the plain at Rexton, southeast, in Mackinac County, as determined by Leverett. The cause for this difference in shore levels is not determined but may have been due to the lodgment, in the McMillan straits, of a large block of stagnant ice which was massed up against the slopes of the moraine. The higher water, in the basin to the east, then must have had its shore pressed against the ice mass, and no records of wave action in this stage were preserved. However, when the water plane had been lowered to the level of 870 feet, the ice mass had either disappeared entirely or had left the slopes of the moraine to allow shore work to become effective there.

Two miles south of Dollarville, the main highway, M-28, ascends an abrupt shore slope to a level, water washed morainic plain which stands at an altitude of 870 feet above sea level. This plain fits the level of the highest shore work at McMillan, which means that the stagnant ice block must have filled completely the entire narrow passage from McMillan to Dollarville. Two small remnants of a higher moraine, one located in the southeast corner of Section 3, and the other just east of the main highway in the west-central part of Section 10, T. 45 N., R. 10 W., stand above the water-washed plain. These morainic fragments reach a maximum altitude of 910 feet and are faced with definite wave-cut cliffs 12 to 20 feet high. (Plate XI, Fig. 1 and 2)

A mile south of Newberry, in the vicinity of the Fire Tower hill, is a fragment of a shore feature at 875 feet. It may be traced as a well

defined shore cliff, from near the tower southeastward for a distance of slightly more than a mile, to the east-central part of Section 1, T. 45 N., R. 10 W. This shore, which surmounts the crest of the moraine, likewise fits the highest Algonquin plane at McMillan.

No other features of sufficient altitude to correlate with the highest Algonquin shore were located south of the Tahquamenon swamp in Luce County. This may be accounted for by the fact that no other areas in the Newberry morainic system reach an altitude sufficiently high to have risen above the water of the lake. Thus the greater portion of the moraine, except near McMillan was inundated and extensively washed by waters of Lake Algonquin when its plane reached the altitude of 900 feet, as recorded at Rexton.

LOWER ALGONQUIN SHORES IN THE TAHQUAMENON REGION

Further withdrawal of the ice sheet from the basin of Lake Superior, coupled with the effects of differential uplift and the opening of new lower outlets, forced a recession in the waters of Lake Algonquin. The records of this recession are preserved in a series of four wave-cut shore terraces which stand between the highest Algonquin and the much lower non-glacial Nipissing shores, in Luce County.

The outermost of these lower Algonquin shore lines stands at an elevation between 795 and 805 feet and marks the margin of the sandy lake plain along the inner edge of the Newberry moraine. It extends from the center of Section 10, T. 45 N., R. 12 W., just two and one-half miles south of Danaher, along the reentrant west margin of the moraine, to the bluff just north of Danaher. Thence it continues eastward along the inner slope of the moraine to McMillan. At this point it swings slightly south and forms the confining wall of the large embayment which continues to the vicinity of Newberry. From this point it continues southeastward for a distance of 6 miles into the northwestern corner of Section 15, T. 35 N., R. 9 W., where it disappears into the swamp. Through much of its extent, the shore line is poorly defined and difficult to trace. The eastern end, south of Newberry, however, is marked by a strongly developed wave-cut terrace which is flanked by a forward strand of cobbly and gravelly material. The shore wash at the base of the cliff contains a large amount of local limestone and dolomite. The more shaly limestone slabs carry numerous fragments of trilobites of the genus *Ogygites*, which Ehlers (32) has ascribed to the Collingwood formation. It is quite probable that this formation lies very close to this locality although no exposures are known because of the complete covering of drift.

Another shore feature, with an altitude ranging between 795 and 810

feet, borders the north or inner slope of the Munising morainic system. It stands as a well marked wave-cut cliff and may be traced as a continuous feature from the northwest corner of Section 10, T. 48 N., R. 15 W., in Alger County for a distance of about 36 miles eastward into the southwest corner of Section 5, T. 47 N., R. 10 W., in Luce County. At this point it breaks into the great Tahquamenon swamp and loses its identity as a definite ridge. The relative position of the shore in the swamp may be traced roughly along a fairly continuous row of well developed sand dune ridges which trend southeastward for a distance of about 14 miles into the valley of the Tahquamenon River near its confluence with Gimlet Creek. These dunes have their steep lee slopes on the south and definitely mark the position of the strand at the time of their formation. They tie up very markedly with the base of the shore cliff immediately south of Cold Spring.

In the easternmost 5 miles of the shore cliff there is a great deal of Hermansville limestone incorporated as slabs in the shore material. This marks the northermost occurrence of calcareous rock in Luce County and may prove of geological significance in working out the contact of the Cambrian-Ozarkian formations, which throughout the area lie so completely concealed.

The cliff-like character of the shore seems to indicate that wave action continued for a relatively long interval while the waters of the receding Lake Algonquin lingered at this level. The sandy and gravelly bars, perched rather high upon the morainic slopes, were formed by storm waves and substantiate the view that deposition accompanied erosion in the shore processes.

The two shore features described above are too high to be correlated with the Lower Algonquin (Battlefield) beaches, which stand at an altitude of 720 feet at Mackinac and 760 to 790 feet at Grand Marais. They are undoubtedly the records of the lower members of the upper Algonquin group which stand slightly above the plane of the (Battlefield) beaches, and were formed during the stage when the Kirkfield outlet was discharging the drainage of the lake.

A further drop in the level of the Lake Algonquin waters or uplift in the basin is recorded by the shore, which follows along the inner edge of the Crisp Point moraine at an altitude of 715 feet. This feature may be traced from the juncture of the moraine with the Sucker basin, in Section 12, T. 49 N., R. 12 W., for a distance of ten miles eastward. Here the morainic segment terminates, and for the next ten miles the shore feature is obscure and may be followed only by level readings. It becomes definite again on the inner slope of the moraine in the northeast corner of Luce County where it can be followed to the vicinity of Crisp Point. This shore pertains to the stage in the Lake Algonquin

PLATE XII




Figure 1.—Algonquin terrace and shore cliff. Nipissing scarp in extreme left. Near Grand Marais.



Figure 2.—Nipissing shore scarp and terrace. Near Grand Marais.

history when the Port Huron-Chicago outlet was functioning and may correlate with the beach at Grand Marais which stands at 725 feet and to which Leverett and Taylor have assigned the name Battlefield.

THE LOWEST ALGONQUIN SHORE

The innermost and lowest shore which may be definitely assigned to Lake Algonquin, in Luce County, parallels Lake Superior at a distance of from one to two miles south of its shore. It may be traced as a definite ridge from the west side of T. 49 N., R. 9 W., westward to the Sucker Basin. Here it swings to the south, follows the escarpment wall around the basin and terminates on the west side of the County. The plane of this shore stands at an altitude of 670 feet and correlates with the beach which has an altitude of 674 feet above sea level at Grand Marais.

During this stage of lake history the basin which now holds the Blind and Dead Sucker rivers was a deep embayment along the shore. This bay was filled with water until the advent of Lake Nipissing, when it was drained.

OLD SHORES IN THE GRAND MARAIS REGION

One of the most remarkable successions of abandoned lake shores to be found on the south Superior shore is located in the region of Grand Marais, in Alger County. In a distance of $2\frac{1}{2}$ miles south of the lake, the terrain rises in the crest of the Munising morainic system, to an altitude of 900 feet above sea level. Here are preserved a series of 10 distinct beach fragments, ranging from the highest Algonquin at 895 feet to the Nipissing, at 625 feet above sea level, mentioned also by Leverett (33).

The highest Algonquin shore is built against the crest of the moraine and is succeeded down slope by several other Algonquin beaches, the lowest of which stands at 820 feet. (Plate XII, Fig. 1) Three beaches at levels of 785, 760 and 720 feet, respectively, representing the lower group of Algonquin shores, are fairly well marked and may be traced for several miles. A well defined shore at an altitude of 670 feet may be followed for some little distance through the area. It represents the lowest level of Lake Algonquin waters and is commonly referred to as the Fort Brady shore although this designation is questionable. With the exception of the three intermediate Algonquin shores, which are represented merely as depositional bars, the features are characterized by wave-cut cliffs which have been carved out of the moraine. Each of the various prominent shores carries a great deal of glacial till and large erratics. (Plate XII, Fig. 2)

PLATE XIII



Figure 1.—Algonquin shore cliff. Near headwaters of Fox River.
S. W. $\frac{1}{4}$, Sec. 2, T. 47 N., R. 15 W.



Figure 2.—Algonquin shore scarp at entry with Fox River Valley.
 $4\frac{1}{2}$ miles north of Seney.

This series of lake shores presents a fairly good picture of the stages in lake succession as melting of the ice and differential uplifts progressively opened new and lower outlets.

ALGONQUIN SHORES IN THE MANISTIQUE BASIN

A prominent shore ridge related to Lake Algonquin activity, and marking a water plane at 870 feet, has been modeled out of the outwash plain in the headwater region of the Fox River, T. 47 N., Rs. 13 and 14 W., in Schoolcraft County. (Plate XIII, Fig. 1). The several headwater branches of the Fox have cut the shore line quite deeply back into the plain so that the scarp now borders the scalloped margin of the outwash in a series of irregular loops. This feature can be traced somewhat interruptedly along the south edge of the sandy plain, to the river escarpment of the East Branch of the Fox River, where it crosses highway M-77, 4½ miles north of Seney. (Plate XIII, Fig. 2)

The rock plateau west of Indian Lake, upon which Cook's moraine is situated, is bordered on all sides by relatively steep escarpment walls which represent the shore cliffs of Lake Algonquin. The elevation of the adjoining plain seems to indicate that erosional activity which carved the cliffs was performed during the highest stage when the Algonquin waters had attained a level of 770 feet. In this, the highest stage of Lake Algonquin, the entire plateau except the low marginal rim, was above the water and stood as an island in the surrounding lake.

In the morainic area east of the Manistique River in T. 42 N., R. 14 W., the glacial drift is very shallow and numerous rock ridges are exposed. This area appears to have been scoured very effectively by wave action of Lake Algonquin. Ribs of limestone and water carved cliffs are prominent in the higher areas. (Plate XIV, Figs. 1 and 2) The irregular surface was examined very thoroughly for evidences of former shores but no definitely connected features were in evidence. It was observed in several of the localities, notably on the eastern slope of the moraine, that the rock cliffs were notched at 700 feet. A gravel ridge in the S. E. ¼, Section 4, another in the center of the same section, are also at 700 feet altitude. (Plate XV, Figs. 1 and 2). Thus it seems probable that the Algonquin waters lingered at this level for a short period, but not long enough to produce very distinctive shore features. Another notch at 670 feet, along the eastern ridge in Section 3, may represent a still lower stage in the recession of the Algonquin waters.

The apparent absence of Algonquin shore lines in the Manistique Basin is due, in part, to the fact that there were very few features of sufficient elevation in the area to have recorded the higher stages. Subsequent scour and erosion by the waters of the Manistique River may

PLATE XIV



Figure 1. A view into Sturgeon Hole, a portion of the Manistique swamp, through the rock corridor in the S. $\frac{1}{2}$, Sec. 14, T. 42 N., R. 16 W.



Figure 2.—Limestone block pavement on crest of a water washed moraine. Most of the till has been removed by water activity. N. E. $\frac{1}{4}$, N. W. $\frac{1}{4}$, Sec. 24, T. 42 N., R. 15 W.

PLATE XV



Figure 1.—Section through a cobbly, gravelly shore showing stratified layers. S. E. $\frac{1}{4}$, N. W. $\frac{1}{4}$, Sec. 4, T. 42 N., R. 14 W.



Figure 2.—A closer view of the feature showing the coarse assortment of material, mostly limestone.

have erased all evidences of shore action against the bordering moraine. Finally, it is probable that recession of the water in the lake was so rapid after the Port Huron outlet was opened, that the various stages in level could not be recorded along the shores.

THE NIPISSING GREAT LAKES

As has previously been mentioned, the Nipissing Great Lakes came into existence when the ice had withdrawn from the valley of the Mattawa-Ottawa rivers and allowed the opening of the outlet at North Bay. "During this stage, the discharge was through the Mattawa River to the Ottawa and thence to an arm of the sea in the St. Lawrence Valley."—Leverett (34). As a result of differential uplift which followed, the North Bay outlet was elevated sufficiently high to raise the water to the level of the St. Clair outlet at Port Huron. At the time when both outlets were in use there was a considerable amount of shore work in progress. The results of this wave activity are reflected in the prominent wave-cut cliffs and strong gravel bars which characterize the Nipissing beach throughout its extent.

The Nipissing Shore Features

LAKE SUPERIOR REGION

The shore of Lake Nipissing in Luce County follows rather closely the present shore, and stands at an altitude of 640 to 645 feet above sea level, or approximately 40 to 45 feet above the level of Lake Superior. The features of this shore are more strongly developed and much more continuous than are those of the Algonquin group. A sharp, well defined escarpment marks the shore for most of its extent in Luce County. Gravelly bars are present on the gentle slopes of the escarpment but wave-cut cliffs and terraces are prominent where the slopes are steeper.

In Alger County, the Nipissing shore forms a prominent escarpment in the village of Grand Marais. It may be traced from the northwest corner of the village, around the south side of West Bay to east-central Section 5, T. 49 N., R. 13 W. Here it swings southward in a rather broad embayment in Sections 4, 9, 10, and 11. In the south central part of Section 11, it turns northward again and leaves the County in the northeast corner of Section 12 at the point where the Sucker River turns westward into Alger County. For much of this distance of 7 miles, the shore is carved out of Cambrian sandstone which stands in vertical cliffs facing Lake Superior. The lower course of the Sucker River, where it swings through the Nipissing plain, was determined largely by the rock wall to its south.

PLATE XVI



Figure 1.—Nipissing terrace with shore cliff in background.
Two miles east of Manistique.



Figure 2.—Sand dunes on Nipissing plain. Ponded slough of
Manistique River in background.

The city of Munising, has been built upon a Nipissing terrace at 640 feet in front of the high moraines whose slopes were washed by the waters of Lake Algonquin.

One of the most interesting and significant Nipissing plains is that in which is situated Lake Au Train. This lake is the remnant of a large Nipissing embayment which was cut off from the larger lake by the construction of a series of bars during the process of lowering of the Nipissing level. No less than sixty such bars, ranging from a few feet to a maximum of ten feet in elevation, are encountered in the mile and a half of road between the head of Au Train Lake and the junction of highway M-94, at the village of Au Train. These bars have been modified since their formation by aeolian activity. The wind, blowing over the sandy plain, has carried sand across the surface and deposited it upon the numerous bars. Secondary sand dune ridges have thus been established on the older dunes, with the resultant development of an irregular surface composed of small corrugations. This area is similar in surface structure to the corrugated feature, in the southeastern corner of Schoolcraft County, south of Milakokia River, which has been described in another section of the report. The corrugations in the Au Train plain are, however, of much lower relief.

The Nipissing Lake along the Superior shore had little or no influence upon the development of the Tahquamenon drainage area. Its waters were confined by heights of land which prevented influx to the marshy lowland tracts.

THE LAKE MICHIGAN REGION

The most prominent Nipissing shore on the Lake Michigan side of the Manistique basin may be observed in the area just two miles east of Manistique. At this point, the main highway, US-2, ascends the sharp shore cliff and emerges on a sandy lake bench 60 feet above the level of the Nipissing plain. The Nipissing shore scarp may be picked up as a definite feature in the southwestern corner of Section 32, T. 42 N., R. 15 W., a half mile east of the Manistique River. (Plate XVI, Fig. 1). From this entry, it follows southward to the Soo line track in Section 9. Here it swings due east and parallels the railroad, on the north side, for a distance of fully 2 miles. In the west-central part of Section 11, T. 41 N., R. 15 W., it turns to the northeast and terminates in Section 1, at the Marblehead Spur, a half mile north of the main line.

Throughout its entire extent, this shore is characterized by an abrupt escarpment built up against a higher sandy lake plain. Where highway US-2 intersects the shore cliff, the sandy fill has a thickness of approximately 100 feet. On the east end of the cliff, where it contacts a morainic

PLATE XVII



Figure 1.—Wave-cut Nipissing rock cliff. City of Manistique.



Figure 2.—Nipissing terrace and wave-cut shore cliff. Three-quarters mile south of Thompson. Sec. 6, T. 41 N., R. 16 W.

slope, the drift is less than 25 feet deep, according to well data obtained by Poindexter.

A spit upon which the city of Manistique stands, was built up from the southwest angle of the shore cliff. It stands at an altitude of 625 feet above sea level or 40 feet above Lake Michigan, which correlates with the Nipissing plane. This spit with its superposed Nipissing dunes has had a considerable influence on the lower course of the Manistique River, as has been mentioned elsewhere in the report. (Plate XVI, Fig. 2)

A wave-cut, rock cliff, marking the position of the Nipissing shore, finds expression within the limits of the city of Manistique. (Plate XVII, Fig. 1). Emerging from the lowland in the northeast corner of Section 8, on the north edge of the city, it trends southward around the west end of the abandoned quarry, to the intersection of Main and Lake Streets. Thence it breaks across to the corner of Terrace Avenue and Cherry Street, paralleling the latter to Steuben Street, where it swings north past the east edge of the cemetery, into the sand dune bluffs. The city of Manistique is located on a Nipissing terrace which slopes gently southward to the lake shore. Nipissing wave action was especially strong on this plain as is evidenced by the extreme thinness of the soil. Bed rock, where not exposed, may be encountered in almost any part of the plain within a depth of four feet.

A Nipissing cliff extends around the rocky headland of Point aux Barques in the southwest corner of Schoolcraft County. It may be followed as a fairly continuous feature northward to the village of Thompson, thence to the south end of the dune which skirts the southeastern corner of Indian Lake. This shore cliff is fronted by a complex of parallel, curving sand dune ridges, separated from each other by narrow, intervening, marshy lagoons. (Plate XVII, Fig. 2)

During the highest stage of Lake Nipissing, the water extended back into large embayments now represented by a number of lakes which evidently became cut off by bars thrown across the headlands at this time. Indian Lake, at 615 feet, Gulliver and McDonald Lakes at 616 feet above sea level are the remnants of the Nipissing embayments. Owing to the broken character of the upland areas, and the extensive marshy tracts surrounding these lakes, definite Nipissing shore features have not been preserved to any great extent, except just north of the station at White-dale where they are difficult to trace out.

POST-NIPISSING SHORE FEATURES

Shore features, associated with a water plane standing between the Nipissing and present Lake Michigan levels, are prominent in certain areas along the shore in Schoolcraft County. The strongest of these shores

PLATE XVIII



Figure 1.—Sandy lake plain on crest of the Nipissing escarpment, two miles east of Manistique. S. W. $\frac{1}{4}$ Sec. 3, T. 42 N., R. 16 W.



Figure 2.—Algoma shore ridge exposing rubble structure. Manistique.

is developed at an altitude of 600 feet above sea level, or at an elevation of 19 feet above Lake Michigan. Its components occur, in places, as wave-cut cliffs along the shore but more notably as rubble bars upon the Nipissing terraces. From information obtained by Hobbs (35), as a result of levelling on the Garden Peninsula, to the south, it is evident that the pronounced shores at 600 feet are correlated with the Algoma stage of lake history.

The Algoma Shore Features

The Algoma shore is not traceable as a continuous feature in Schoolcraft County, but occurs rather as more or less disconnected units sometimes widely separated, along the shore. In certain areas, the Algoma features are very definite but in the main they are either obscured by low dune developments, or missing entirely as a consequence of later wave activity.

Less than a half mile south of the village of Thompson, fronting the Nipissing cliff, is a small trace of the Algoma shore. It extends for a distance of about three-quarters of a mile as a wave-cut scarp, along the west side of Section 6. A series of fore-dune ridges have built up in front of the low cliff and extend outward to the present shore.

A group of three parallel rubble beaches, with more or less abrupt front slopes, was built up on the Nipissing plain in front of the shore-cliff, on the west side of the city of Manistique. The outermost of these bars stands at an altitude of 600 feet above sea level and has been ascribed to the Algoma shore by Miss Stevenson (36). (Plate XVIII, Fig. 1). The two higher beaches are transitional between the Nipissing and Algoma levels and were formed either during the lowering of the Nipissing waters or by storm waves during the Algoma stage. A great deal of rubble has been piled up on the Manistique headland in a series of parallel ridges at levels between the Algoma and the present beaches. Much of the rubble rests directly upon the limestone floor and is composed of material which has been broken from it by the waves. The ridges are low, but usually well developed. They may possibly represent storm wave deposits of the present lake.

Miss Stevenson (37) mentions a very definite cliff against the front slopes of the dunes on the Nipissing spit east of Manistique. She ascribes this cliff to the shore work of Lake Algoma on the basis of certain facts which are very obvious. She states further that, "the termination of the Nipissing sand dunes in the Algoma cliff represents the time when the down-cutting of the outlet, the uplift, or both, were halted and erosion by the waves became dominant in this area." This conclusion seems justified on the basis of the extensive stretches along the shore which have

PLATE XIX



Figure 1.—Gently dipping limestone floor along shore of Lake Michigan.
South of Seul Choix Point.



Figure 2.—Glacial erratics on limestone lake floor. Along shore of
Lake Michigan just east of Manistique.

been scoured clean by the waves, to expose the limestone pavement. (Plate XIX, Fig. 1). The numerous glacial boulders and erratics distributed on the rock floor are merely the concentrated remains of till features which have been completely destroyed. (Plate XIX, Fig. 2). The fore dune ridges, gravelly bars, and sandy beaches which have been built up on the rocky platform between the Algoma cliff and the present lake, support her view that "since Algoma time deposition has been more important than erosion."

There are, no doubt, many other areas along the shore where Algoma features could be definitely worked out, but very detailed work such as carried on by Scott, Stevenson, and others will be necessary to accomplish such determinations.

Chapter V.

AEOLIAN ACTIVITY

One of the most significant physiographic aspects of the Tahquamenon-Manistique drainage basins is related to the activity of the wind which was effective during glacial as well as post-glacial times. Wind-developed sand ridges and dunes comprise a relatively large proportion of the low lying marshy tracts of the region, and frequently are responsible for such complications as to make differentiation of the features very difficult, if not impossible.

The dunes of this area, may for the sake of convenience, be subdivided into three major groups, as follows:

- (I) Shore Dunes along Lake Superior.
- (II) Dunes of the poorly drained marshy, lowland plains of the main drainage ways.
- (III) Shore dunes along Lake Michigan.

These will be discussed more or less in detail but special emphasis will be placed upon those of the second group.

I. Shore Dunes Along Lake Superior

Dunes of variable extent occur intermittently along the shore of Lake Superior in Alger and Luce counties and, in certain localities are of especial geologic significance.

A. ALGER COUNTY.

(Shore Dunes)

The sand dunes in Alger County are concentrated principally along the present shore of Lake Superior. Smaller dunes are distributed here and there on the shores of older lakes and occasionally on the margins of the larger swamps and marshes but are of little importance.

The most important, and by far the greatest development of sand dunes is found along the shore in the region of Grand Sable Lake in the vicinity of Grand Marais, in the northeastern part of the County. The Grand Sable dunes in this region represent, without question, the most imposing mass of wind deposited sand on the entire south shore of Lake Superior. They cover in total an area of approximately five square miles, and possess crests which attain a maximum altitude of 957 feet, or an elevation of 355 feet above the level of Lake Superior.

PLATE XX



Figure 1.—Crest of Grand Sable dunes showing deflation hollows.
Near Grand Marais.



Figure 2.—Exhumed forest on crest of wind-swept sand dunes.
Near Grand Marais. Lake Superior in background.

The topography of this dune area is especially rough and broken. Deep hollows and blowouts, extending at times to the original floor, have been scoured by wind deflation. (Plate XX, Fig. 1). The deeper bottoms of the blowouts in the area between Grand Sable Lake and Lake Superior are floored locally with small patches of coniferous growth suggesting a water table not far below. In the shallower bottoms, vegetation is scant and entirely lacking especially where the strong northwest on-shore winds sweep through. In places, along the more sheltered slopes, leguminous plants have been successful in gaining a foothold, and through their roots serve to bind the sand somewhat more firmly together. Exhumed forests on the wind swept crests of the dunes are evidence of post-dune wind activity. (Plate XX, Fig. 2)

On the exposed slopes, along the shore of Lake Superior, where a section may be obtained from the lake level to the crest of the dunes, many interesting features are revealed. The exposure portrays an aeolian character only near the surface. At the base of the shore cliff, boulder-clay of glacial origin extends upward 100 feet or more above the level of Lake Superior. The till is a mixture of sand, silt and clay, beset with numerous erratics of large dimension. Resting upon this glacial floor are a series of stratified beds consisting of an assortment of rounded cobbles, coarse gravels, sands and silts which continue upward to an elevation of approximately 280 feet above the lake level. The low sand dunes, along the shore, are perched upon this lacustrine formation at an altitude of 870 to 880 feet. (Plate XXI, Fig. 1)

From the juxtaposition of the stratified material in this section it would appear that the sand dunes, especially those along the shore, are situated on an older lake feature, possibly a bay-mouth bar, which had been thrown across the head of the embayment to form Grand Sable Lake, sometime during early Algonquin times. This suggestion is merely speculative and will require more detailed studies for conclusive proof.

The general absence of a vegetative cover in the Grand Sable section of the dunes has been responsible for a considerable amount of translocation by the wind. Immediately northeast of the lake, a forest is now in the process of being invaded by drifting sand. The sand has piled up to such a depth as to completely cover the smaller trees, and the larger ones are buried to half their total heights. The channel of Sable Creek, the outlet of Grand Sable Lake, is being continually shifted eastward by the rapidly invading sand. (Plate XXI, Fig. 2) Farther west, the sand is gradually encroaching upon the margin of Grand Sable Lake, and much material from the dune is being carried into its water.

In the deflation hollows or blowouts, the atmosphere is set into a turbulent motion as a consequence of the numerous cross currents and eddies which are developed by the lashing gales of Lake Superior. The

PLATE XXI



Figure 1.—Perched dunes resting on cobbly bar 280 feet above level of Lake Superior. Grand Sable area.



Figure 2.—Sand dunes encroaching upon channel of Sable Creek. Near Grand Marais.

PLATE XXII



Figure 1.—A lag-gravel plain with perched sand dunes in background.
Grand Sable dune area, near Grand Marais.



Figure 2.—Mud streams on north flank of Grand Sable dunes,
near Grand Marais.

effect of this turbulence is largely that of sorting out and removing the finer particles and concentrating the coarser material to form a desert armor. Ripples are likewise produced and vary, according to texture of the material included and exposure to shifting wind currents, from minor rippled forms to giant waves, as mentioned by Bucher (38). One of the largest giant ripples observed after a severe wind, measured seven feet from crest to crest and had a height of 14 inches. This ripple contained faceted and polished rock fragments ranging from coarse gravel to angular pebbles of over an inch in diameter. In general, it may be stated that the ordinary ripples are for the most part veneered with a thin covering of coarse sand and fine gravel which acts in the capacity of preserving the constructional forms. It seems to follow also, that the size of the ripple is a function of assortment of particles and wind velocity, for the larger forms always contain the coarser material.

Along the Superior shore, where the perched dunes are relatively low and the sand mass is limited, the wind has successfully removed the bulk of the finer debris with the result that in local areas there has accumulated a concentration of coarse material on the residual floor to which Udden (39) applies the term, lag gravel. (Plate XXII, Fig. 1) Where complete removal of the dune sand has been accomplished, the cobbly material of the former beach is exposed.

On the less sheltered areas, where the finer particles have been removed, the long continued sand blasts have played upon the surface pebbles and cobbles to develop various types of wind-worn stones, to which Bryan (40) has collectively assigned the term "ventifacts". Many of these stones have been faceted by wind driven sand blasts to produce the characteristic *Ein, und Drei-kanter* of the German literature.

In the dune region west of Au Sable Lake, the forms attain their maximum development with crests reaching an altitude of 957 feet. This sand covered area is densely forested with a growth of hardwood which affords protection against deflation. Except in local areas, where the timber has been removed from the slopes, the sand is stable and very little shifting by wind has occurred.

In view of the geological evidence at hand, it seems reasonable to conclude that the Grand Sable dunes were formed from contributions of sand of a higher shore, rather than of sand from the present beach. In the first place; the dunes are perched upon a shore which stands at an elevation of 280 feet above Lake Superior. This shore is too high to correlate with the Nipissing and is undoubtedly related to one of the higher Algonquin lake levels. Secondly, the wave cut cliff which fronts Lake Superior is too high and steep walled to have allowed much movement from the present shore. Thirdly, the present beach, for most of its extent along the sand dune area is composed essentially of boulders,

cobble and gravel and scarcely any sand is now being reworked by the waves. Hence, little, if any, material of the present beach is fine enough in texture to be transported inland very far, even by the strongest winds blowing off Lake Superior.

The Grand Sable Bluffs

The Grand Sable dunes terminate in the central portion of Sec. 7, T. 49 N., R. 14 W. At this point and westward, for a distance of about a mile, the dunes give way to the steeply sloping Grand Sable banks. (Plate XXIII, Fig. 1) Aneroid measurements show these bluffs to rise to an altitude of 855 feet or 253 feet above the level of Lake Superior. The bluffs are composed essentially of water rounded quartz sand intercalated with lenses of gravel and pebbles of varying texture and composition. They slope down to Lake Superior with an average angle of repose of 35 degrees. Much sandstone, of local derivation, both rounded and in blocks, as well as numerous water-worn erratics, are included. Near the base of the bluff there is considerable silt, with the character of glacial till, which forms the matrix for the glacial boulders. The crest of the bluff is a wind-modified sandy lake plain over the surface of which small dunes have been developed. The sandy-gravelly bluffs terminate in Sec. 12, T. 49 N., R. 15 W., where the rocky headland of Cambrian sandstone sets in to skirt the shore around Au Sable Point.

Another prominent area of dune development lies in the region east of East Bay, near Grand Marais. This region of dunes continues along the Superior shore and extends for a considerable distance eastward into Luce County. The dunes in this section are definitely aligned in more or less paralld ridges, separated by intervening longitudinal depressions. They are not perched, but rather have their bases at the shore level of Lake Superior. Thus, their summits are lower than those of the Grand Sable section and range from 40 to 60 feet above the level of the lake. These dunes are heavily forested, or otherwise blanketed with vegetation, and the sand in them is fixed, except near the immediate shore where small fore-dune ridges are in process of formation.

A limited area of dune features has been developed along the north side of the Sucker River in the south eastern part of T. 49 N., R. 13 W., and the north eastern part of T. 48 N., R. 13 W. These dunes have a general trend from northwest to southeast and extend more or less parallel to the Algonquin shore. They are arranged in definite ridges but have a relief rarely exceeding 30 to 40 feet. (Plate XXIII, Fig. 2)

In the region north of the Beaver Lakes, low piles of sand have been heaped up by the wind. They are more or less barren of vegetation, and subject to much movement by drifting. The windward slopes of these dunes lie some distance back from Lake Superior but the back slopes

PLATE XXIII



Figure 1.—Sand bluffs near Au Sable Point.



Figure 2.—Sand dunes in the Sucker River basin.
Sec. 26, T. 49 N., R. 13 W.

have been carried leeward to the north shore of the Beaver Lakes, to form marginal features. This dune section varies in width from less than an eighth of a mile on the southwest end of Little Beaver Lake, to slightly more than half a mile on the west side of the Beaver Lake outlet.

A broken line of narrow dunes extends along the Superior shore, around the Grand Portal headland from Chapel Beach, for a distance of seven miles to the westward. These dunes are relatively low and lie directly behind a rocky shore. They were formed at a higher stage of lake level when a sandy beach was being developed along the margin. Except in the coves and reentrants, practically no sand is being contributed to these dunes by the present beaches.

A series of low, but rather well formed dunes have developed along the mouth of Au Train River just to the north of the village of Au Train. This sandy tract extends for a mile along the shore of Lake Superior and has an average width of about six or seven hundred feet. Much of the material in these dunes has been brought in by the sluggish Au Train River, which in its course from Lake Au Train, meanders broadly across the low-lying sandy Nipissing plain. Lake Superior is likewise contributing new material to the growth of these dunes which in most of their extent are barren of vegetation and subject to much movement by the wind.

B. LUCE COUNTY

(Shore Dunes)

There are three definite areas of dune development along the shore of Lake Superior, in Luce County. The most westerly dune lies north of the Sucker Basin in Twp. 49 and 50 N., R. 12 W. and is a continuation of the sandy ridges which may be traced eastward from East Bay in Alger County. Unlike their counterparts on the west side of the Sucker basin, these dunes are relatively high and attain crests which rise to 100 feet above Lake Superior. They are clothed with vegetation, and except for the fore-dune ridges along the shore, are quite definitely fixed. The position of these dunes, situated against the wave cut Nipissing bluff, would suggest a post-Nipissing origin.

An area of dunes along the lake shore in R. 9 W. has been built up into a series of low tenuous ridges on a post-Nipissing plain, from material of the present beach. They are well clothed with vegetation and thus lie protected from serious translocation by the wind. The bulk of the aeolian sand in this area lies north of the Two Hearted river and appears to terminate near its mouth. Fragmentary patches of drifting sand mostly in the form of fore-dune ridges on the present beach, may be picked up here and there, eastward to Temple Lake.

The most easterly dunes in Luce County are situated along the shore of

Lake Superior in Twp. 8 N., where they may be traced to Crisp Point on the county line. They stand upon the Nipissing terrace and, in all probability, were blown up by wind from the Nipissing strand.

II. Dunes of the Poorly Drained, Marshy, Lowland Plains of the Main Drainage Ways

The most extensive single physiographic feature of the area included in this report is the marshy lowland tract associated with the drainage basins of the Tahquamenon-Manistique rivers in Luce and Schoolcraft counties. This lowland, marshy plain represents the floor or bed of glacial Lake Algonquin, the predecessor of Lake Superior. The plain, as a whole, is poorly drained and, since its occupation by the Algonquin waters, has received a vast accumulation of organic matter which is embodied in a shallow mantle of muck and peat resting upon the lacustrine sand beneath. (Plate XXV, Fig. 1)

The level monotony of this extensive drainage area is broken by innumerable ridges and knolls of sand, which Scott agrees, are deposits of aeolian activity and may definitely be classified as dunes. Thousands of these dunes are distributed through the lowland tract and range in size from merely a few feet to 60 and more feet in height, and from a few hundred feet to several miles in linear extent. In a broad way, they follow the mainly developed drainage channels of the region, but are also aligned as east-west ridges along the successive shores related to the retreatal fronts of Lake Algonquin. Scott has suggested that these continuous rows of dunes mark the position of former shores and represent definite shore features.¹ The dunes of this area were formed before the organic deposits were developed at a time when the dessicated sand in the broad, shallow, river flats and along the lake shores was fully exposed to winds, and moved by them. Because of the poorly developed drainage facilities of the area, the water table has always stood rather high and its position has limited definitely the depth to which translocation, as described by Free (41) could develop.

The building up of dune successions, especially along the borders of the receding lake, was responsible for considerable derangement in the drainage lines. Sand ridges, thrown as barriers against the streams, which were then developing in the wake of the lowering Lake Algonquin, caused a damming of the youthful drainage and a consequent impounding of the water. Thus the streams were forced to overflow their shallow channels and spread out to form the marshy interfluves.

Many of the dunes, particularly those which follow the drainage courses, are irregular in shape but relatively symmetrical in profile.

¹Scott, I. D., Personal communication in the field.

Those forms, however, which are associated with former beaches and possess a trend from east to west are usually characterized by gentle windward and steeper leeward slopes.

The divide between the waters of the Tahquamenon and Manistique rivers roughly defines the line along which the Algonquin waters split when recession was in progress. North of the divide, in the Tahquamenon basin, the shore dunes carry their steep leeward slopes, facing southward; but south of the divide in the Manistique basin, dunes have their steep slopes facing northward. This condition of opposed dune slopes in the two basins is significant in that it throws much light on the problem of recessional activity of the waters of Lake Algonquin.

It seems logical to assume that the dunes in this area were formed in much the same manner as post-glacial dunes have formed and are still forming. It is obvious then that the winds which sculptured the aeolian ridges must have been on-shore winds in order to have moved beach material landward. The opposable slopes of the dune ridges thus bear out the conclusion that the Algonquin waters parted in the region of the Danaher divide with the result that two basins ultimately formed. The water in these basins continued to recede, being drained out, in part to the north and in part to the south.

That the lowering of Lake Algonquin was the result of gradual and not rapid recession is manifested by the numerous parallel rows of dunes which lie between the main divide and the present basins of Lake Superior and Michigan. In many of the areas, ridges of sand 50 to 60 feet in height have been piled up along the older shores and present evidence of relatively long intervals of temporary halts in lake recession, during which accumulation of sand was in progress.

In many parts of the Manistique drainage basin, the features are so complex as to make complete differentiation impossible. Ridge after ridge of sand, separated by intervening marshy lanes, and interspersed with small tracts of sandy lake plain, stretch for miles across the lowland region and in places cover entire townships. So complicated is some of this country that the individual features are not revealed, nor can they be separated in the areal mosaics constructed from photographs on a scale of 4 inches to the mile. Such areas have been designated *complexes*.

Two distinctly different lowland complexes may be distinguished in the Manistique river area. One of these has been given the name of Seney complex, from its type development in the vicinity of the town bearing this name. The other has been termed the Manistique complex because of its distribution along the shore of Lake Michigan near the city of Manistique.



Air photo of fore-dune ridges, near Seul Chole Point, Schoolcraft County

THE SENEY COMPLEX

This feature is essentially of an interior character and consists of wind blown sand in the form of ridges and knolls, interspersed with open marshy areas and muskeg. The dunes may be present as isolated forms scattered through broad tracts of lowlands, or they may be connected as more or less continuous and closely spaced ridges separated by intervening marshes, and poorly drained swamps. The sandy uplands make up from 25 to 75 per cent of the complex area and are usually covered with Norway, white and jack pines. The marshy lowlands constitute the remainder of the region and are grown over to grasses, sedges, cassandra and occasional timber size forest trees.

The dunes in the Seney Complex are extremely variable in size and extent. They range from mere waves, a few feet high covering a sandy lake floor, to extensive dunes, 50 to 60 and more feet in height. Some of the knolls are of distinctly local character and occur as individual small patches. The ridges, on the other hand, may continue across country for miles without interruption. Extensive areas of sandy lake floor break into the complex, but are generally much modified and show very definitely the effects of translocation and deposition by the wind.

THE MANISTIQUE COMPLEX

The Manistique complex is concentrated largely along Lake Michigan and consists of a succession of narrow, parallel ridges which follow closely the trace of the present shore. The ridges are separated by narrow swales, seldom exceeding 2 chains in width. They are timbered with Norway, white, and jack pine whereas the associated swales have a cover of spruce, cedar, and tamarack, with cassandra or grass in the open areas.

The dune ridges in the Manistique complex generally follow closely along the present shore, often in the form of fore-dune features. Progressing landward, the ridges become successively higher and higher until finally they merge in the hinter dune area and lose their individuality in the massifs.

The swales are quite uniformly interspaced with the ridges and give to the relief a character which is typically undulating. Because of its structural character, the writer here proposes to describe the relief as "corrugated." A beautiful example of corrugated surface may be viewed in the southeast corner of Schoolcraft County, north of Lake Michigan in the vicinity of the Inland Steel Company harbor. (Plate XXIV)

In the distance of a little more than a mile between the lake shore and Milakokia river, in this area, there are a series of 36 low, parallel sand dune ridges ranging from 5 to 15 feet in height. They are separated from each other by narrow marshy lanes. Starting as low fore-dune

ridges along the shore they become successively higher and more completely developed as the distance from the shore increases. North of the Milakokia river, the ridges give way to a massif of dunes which reaches a maximum elevation of 125 feet above the level of Lake Michigan and terminates completely at the swamp which borders the shores of Shoepack lake.

In the area, between the well defined inner fore dune ridges and the main massif, is a transitional belt of deformed ridges. These ridges are roughly *en-echelon* in arrangement and present their scalloped hollows windward. It appears entirely probable, as Scott has suggested, that the waves on the prograding shore may have initiated the hollows in the deformed dune areas, which later were subjected to wind modification.

The fore-dune ridges, built up in succession along the receding shore are moderately symmetrical in shape. They do not possess the characteristic leeward-windward slopes, and for the most part the profile is quite uniform. There seems good reason to believe that the parallel ridges may possibly correspond to off-shore lacustrine bars which have more recently been modified by aeolian activity.

Opportunity for study of a small isolated dune section, associated with the marshy tract, is afforded in the vicinity of the fire tower located a short distance south and east of the village of Seney. The tower is situated on the crest of one of the highest dunes in the region and overlooks a vast expanse of open muskeg and timbered swamp on all sides. The dunes here are well developed and possess the characteristic leeward and windward slopes. They rise to a maximum altitude of 800 feet or to an elevation of 60 feet above the surrounding marshy plain. The dunes are here massed together and rather closely crowded into a compact area of only a few square miles. (Plate XXV, Fig. 2)

The features are somewhat complicated in structure and present a variety of forms. Piles of sand without trend break off into definite ridges which, with their associated intervening sand-floored basins may be traced uninterruptedly for some distance eastward. Deflation hollows and blowouts are associated with certain portions of the district and occasional forms having the appearance of barchans may be viewed from the tower.

Close investigation of the area, especially in the valleys and the basins, reveals the presence of glacial till and boulders which form the foundation for the dunes. It is very probable that during the interval of dune development a small isolated mass of moraine standing along the margin of Lake Algonquin in its southward retreat from this area, intercepted the sand which was whipped up from the adjacent shore and was in large

PLATE XXV



Figure 1.—A portion of the Manistique swamp. Near Walsh.



Figure 2.—Sand dune topography near Seney as viewed from the Fire Tower—looking northwest.

PLATE XXVI



Figure 1.—Dune blown up against the slope of a moraine in the Manistique swamp. N. E. $\frac{1}{4}$, Sec. 21, T. 42 N., R. 15 W.



Figure 2.—Section of dune showing cross-bedding in the unconsolidated sand. Tahquamenon swamp, two miles north of McMillan. Sec. 21, T. 46 N., R. 11 W.

measure responsible for the aeolic accumulation. Similar features prevail in other areas where morainic ridges border the swampy lowland. (Plate XXVI, Fig. 1)

INLAND DUNES

From detailed studies made of the features in the marshy complex of the Manistique swamp, it is apparent that the majority of the dunes were shaped by the wind along prograding shores of Lake Algonquin. This is especially true of those ridges which have an east-west trend across the lowlands.

On the other hand, however, there are a multitude of aeolian features which appear in no way to have had any association with retreating shores. Dunes of this type are distributed along some of the major drainage ways and follow, in general, the trend of the stream courses. To a large extent, they are most strongly developed on the west banks of the southeastward flowing rivers, a fact which would demand prevailing wind activity from the northeast.

In connection with these dunes, the question arises as to whether they should be classed as shore dunes or strictly as inland types, following the classification of Solger (42), Högbom (43), and others. The writer is inclined to consider them as inland features and suggests an origin which is correlated indirectly with stream activity.

In describing the general physiographic character of the Manistique swamp on pages 59-61, mention was made of the almost universal distribution of shallow bodies of muck and peat over the lake laid sandy floor. Close studies of a large number of dune areas along the rivers failed to reveal a single major dune form which has been developed upon a base of muck or peat. So far as could be determined, sandy plains provide the foundation for the structures. Hence, it follows that the dunes were formed upon flood plain areas in the lowland tracts. As the rivers in their early history eroded their channels into the lake plain, a large amount of sandy material was released. Much of the finer sand was washed up on the flood plains and flats along the river borders during each flood. Exposed during periods of low water, the sand became desiccated and in such a condition was easily moved by the prevailing winds and then piled up into definite ridges along the lee banks.

Again, these dunes may have been developed by transportation and deposition of older deposits of aeolian sand which had previously been laid down along the lake shores. Thus modified, should they still be classed as inland types or would they better fit into the category of shore dunes?

The storm winds in the region are at present from the northwest off Lake Superior, whereas the position of the dunes in the stream areas

call for a prevailing northeast wind. This discrepancy in wind direction need not be an obstacle to the solution of the problem inasmuch as a large mass of ice still persisted to the north and east of the Superior basin. Anticyclonic storms generated over the ice area, may have been responsible for the directional change during Algonquin times, following the theory advanced by Hobbs. (44)

III. Shore Dunes Along Lake Michigan

Sand dunes occur intermittently along the entire shore of Lake Michigan, from Point aux Barques to the Inland Steel Plant, in Schoolcraft County. Throughout most of this distance they border the lake and constitute a complex of ridges which are aligned more or less parallel to the shore. The ridges are separated from each other by interspaced lanes of peat-filled marshes and occasional sandy, lake plain flats.

The dunes in this area are variable in dimension. They range from low, narrow fore-dune ridges now in process of being piled up on the present shore, to rather bulky massifs which rise to 125 feet and more above the level of Lake Michigan, in the southeastern corner of Schoolcraft County. According to studies made by Scott and Miss Stevenson of the Manistique area, there have been three stages of dune activity. These are ascribed to Nipissing, Algoma and present shores. The largest and most deformed dunes of the region have been traced back and correlated with the Nipissing beach which here stands at an altitude of 625 feet, or 40 feet above the level of Lake Michigan. The Algoma dunes can be definitely tied up with a beach at 600 feet. Dunes of this stage are low and under-nourished and in no way comparable in bulk to the Nipissing features. The recent forms, as has already been mentioned, are mainly low, tenuous fore-dune ridges seldom exceeding ten feet in height. They are being constructed from material of the present beach and are evident wherever a supply of sand is available. The slopes of these young ridges are more or less uniform and symmetrical and, although not representing the angle of repose of the included sand, appear to be slightly steeper on the leeward sides.

Dunes are somewhat lacking along the shore in those areas where the floor of the lake is composed of bed rock. These dune-barren localities are situated primarily in the headland portions of Point Aux Barques and Seul Choix Point. Another extensive tract of rocky shore lies between the city of Manistique and the Nipissing shore scarp a few miles to the east. It is significant to note, in the areas where the limestone beds form the lake shelf, that an abundance of large erratics are scattered over the surface. These boulders are the concentrates of former till deposits, the finer material of which has been completely re-

moved by subsequent wave action. It is quite possible that they represent the remnants of glacial features which may have correlated with the final stages of activity in the Lake Michigan lobe. This view, however, cannot be conclusively proved because of the lack of sufficient evidence.

The dunes of the Lake Michigan shore have played an important role in the history of the Tahquamenon-Manistique drainage basins. They were responsible, to a considerable degree, for the damming of the newly formed rivers in Nipissing times. This damming of the drainage in the lower reaches caused flooding in the interior lowlands and resulted in the swampy, poorly drained condition which characterizes the region.

From the fact that the shore dunes of the interior lowland tract are associated with the activities of Lake Algonquin, it seems logical to conclude that drainage was free and open during most of this time. This view is based on the consideration that the sandy lake flats had as yet not been encroached upon by vegetation which was responsible for the accumulation of the vast expanses of shallow muck and peat. Thus, a continuous supply of sand was originally available for wind translocation.

With the close of the Algonquin stage, the waters were lowered to the Nipissing level and remained in this position for a considerable time as is evidenced by the prominent wave-cut cliffs which mark the shore near Manistique. The building of dunes on the newly formed lake terrace, eventually blocked the outlet of the Manistique River, and at the same time likewise affected its tributaries in the hinter plain. It was during this period of flooding, that organic activity became prominent and peat was laid down upon the sandy floor as a protective buffer against further ablation.

When the Manistique River finally opened a new outlet into Lake Michigan the flooded waters were carried out. The tributary streams, with headwater areas uplifted by the differential movements which had been in progress through much of Algonquin times, became actively engaged in cutting channels into the sandy lake floor. This renewed fluvial activity in the tributary streams provided a new source of sand which, blown up from the flats and flood plains, became piled up into dunes along the river courses.

It is thus possible to trace out two major episodes of dune formation in the Tahquamenon-Manistique drainage basin. The earlier of these episodes may be correlated with wind activity along the retreating shores of Lake Algonquin. The dunes of this stage are definitely of the shore type. The later dune developments are tied up with the Nipissing activities along the Lake Michigan shore and the consequent formation

of inland features along the stream courses. An interval of flooding and peat accumulation separates the two successions.

SUMMARY AND CONCLUSIONS

The Tabquamenon-Manistique Drainage Area, representing the most extensive tract of marshy lowland in the state, seems to hold a key position in the final interpretation of the Pleistocene problems of the Belted Lowlands province of the Northern Peninsula. The activities which prepared and ultimately shaped the surface expression of the area are associated with the waning stages of the Labrador ice sheet in late Wisconsin time. Three distinct episodes are represented in the development of the surface features; these are related to glacial, lacustrine and aeolian processes.

Glacial Activity

In the first episode, that of glaciation, it is significant that the region was affected largely by a readvance of the ice from the Superior basin. The movement of this ice mass, as recorded by striae on bed rock, was largely south-southwestward, in contrast to the generally westward direction of an earlier invasion. It is quite evident that the Green Bay ice block which still persisted to the west was responsible to a large degree for the newly directed movement.

Glaciation has left its stamp on the character of the region in the form of two strongly developed morainic systems, which I have named the Newberry and the Munising moraines, respectively. The development of these morainic systems is contemporaneous with the final activities of the Green Bay lobe in the region of Trenary till plain. It seems plausible, from the related positions of the moraines of the Green Bay and Superior lobes, to assume that the ice in the Green Bay lobe had become stagnant before the glacier completely retired from the west end of the Munising moraine.

Throughout the area, no features can definitely be ascribed as having been formed by the Michigan lobe before its final disappearance. There is a possibility, however, that the bouldery concentrates, including large erratics, which fringe the rocky shore of segments of Lake Michigan in Schoolcraft County, are the remains of the final till deposits of the Michigan lobe. Owing to the completely modified character of the surface and the fragmentary bits of evidence at hand, it is possible only to make conjectures concerning such a relationship. Nevertheless, it seems wholly within the province of good judgment to believe that the final position of the Lake Michigan lobe was somewhere between the present lake shore and the front of the Newberry moraine. Possibly the

Michigan lobe retired much in the same way as did the Green Bay lobe, as a stagnant body of ice, and may have been overridden to some extent by the readvancing sheet of ice from the Superior basin.

Although the moraines which were left by the Superior ice lobe are fragmentary and disconnected because of subsequent erosional activity, they were undoubtedly continuous in their trend for some time after the ice retired from them. Hence through the interval when lacustrine activity was paramount, they served as barriers, at least temporarily, against rapid outflow and were thus responsible for much of the ponding of water which occurred in the early development of the basin.

With the exception of a few localized areas where it has been completely fragmented, the Newberry morainic system forms the outer margin of both the Tahquamenon and Manistique drainage basins. This moraine served as a control in basin development for the area and likewise determined definitely the course of the Manistique River and to some extent the course of the lower reaches of the Tahquamenon. If this moraine had not been deposited in the position which it now holds, it is barely possible that neither of the two large river systems could have been initiated. Under such a set of circumstances there would have developed upon the tilted lake plain a series of parallel independent drainage lines such as are now represented by the numerous branches of the Manistique River. Carrying the speculation of such a condition a step further, it seems reasonable to venture that the marshy lowlands would have assumed much more extensive proportions to the east of the present major drainage lines.

Lacustrine Activity

The second episode, related to lacustrine activity, did not begin until the body of the Superior ice sheet had disappeared largely from the surface of Michigan and the Superior basin was freed of ice at least to the meridian of Wetmore. At this stage, the Anna River at Munising carried discharge water of the partially formed Lake Algonquin across the high moraines south of Munising, into the head of the Manistique swamp. The contribution of water through this passage was relatively small and the outlet was short lived as is manifested by the small amount of scour and the narrowness of the valley. The occupation of the valley by the Algonquin waters is evidenced, however by the presence on the slopes of the moraine within the valley, of intercalated lacustrine silts, clays and sands to an altitude of 838 feet above sea level. These deposits are thinly laminated and can no doubt be classed as varves, according to descriptions by Antevs, (45) Reeds (46) and others.

A further retreat of the ice northeastward in the Superior basin, and

the attendant differential uplift in the region, resulted in widespread inundations. The water from Lake Algonquin found its way across the lowland areas of Luce County and spread out in sheets over the interior sandy plains. During the highest stage in lake level, the greater portion of the relief was under water, merely a few highland tracts standing high enough to remain above. In this stage of lake development wave action was effective on the crests and slopes of the higher moraines, especially those that were either slightly above, or submerged in, shallow water. Thus, much of the relief was modified, with the result that slopes were washed down and smoothed out and the original irregular surfaces became flattened to a considerable degree.

An important passage for the waters from the Superior to the Michigan basins was affected through the Tahquamenon depression across the narrow straits at McMillan, into the Manistique lowlands. The divide, near Danaher, stands at an altitude of 720 feet and represents the lowest passage established between the two areas.

The continued uplift of the Superior basin resulted in the diversion of more drainage waters through the Port Huron outlet and caused the plane of Lake Algonquin to be lowered. When the waters were too shallow to cross the divide in the McMillan Straits, they split. Part of the water drained out northward through the Tahquamenon depression, and the remainder found its way southward into the Manistique drainage basin. This parting of the water into two separate water sheds is evidenced by the opposing slopes of the dunes on either side of the divide. The northern dunes have their steep lee slopes facing southward whereas the southern dunes have the steep slopes facing northward.

The Tahquamenon and Manistique rivers did not come into existence as definite water courses until the Lake Algonquin level had been lowered sufficiently to allow channels to be etched into the sandy lake floor. They increased their length, not alone by headward erosion, but largely through extension of the lower reaches as the water in the retreating lake subsided. Throughout the process of river development, uplifts were effectively in progress. The Lake Superior basin, being farther removed from the hinge line, naturally suffered a greater amount of movement than did the Lake Michigan basin. The result of the differential uplift activity was to raise the rock sill of the Tahquamenon River at the Upper Falls to a point where the drainage could no longer overflow. The ponded waters spread out into the low lying flats and a large shallow lake was formed in the basin.

The differential uplift, on the other hand, raised the headwater area of the Manistique River to a much greater degree than its outlet. This caused the river to be rejuvenated, with the result that it was able to deepen its channel materially.

Aeolian Activity

The third episode involved in the Pleistocene history of the Tahquamenon-Manistique drainage area is concerned mainly with aeolian activity. Wind work, on a large scale, was initiated at the time when the waters of Lake Algonquin began to subside in the interior basin and continued well into and through Nipissing times. The majority of dunes within the drainage basins were formed during the retreatal stages of Lake Algonquin. They were blown up into ridge rows from the successive beaches on the prograding shores and represent the various positions of the strand as the water level receded.

Although shore dunes constitute the larger proportion of aeolian features in the region, it is nevertheless significant that extensive areas along the major river courses are beset with dunes. These have been developed on the flood plains and river flats out of material which was translocated by running water. They seem to bear no relationship whatever to the receding shores of the older lakes, but rather parallel the major valleys. Hence, they are not shore dunes in the true sense of the word and because of their inland origin may well be classed as interior dunes.

The development of thousands of dunes scattered broadcast over the lowland area, has been responsible for a great deal of derangement in surface drainage. The dunes which formed along the older shores impounded much water during lake retreat and prevented its complete removal. Lagoons, formed behind the numerous shore ridges, became sites for organic accumulation which is reflected in the almost universal distribution of muck and peat over the sandy lake floor. Broad interstream areas of marsh and muskeg give further testimony to the ponded condition of surface drainage, subsequent to dune formation.

In much of the drainage basins the dune ridges are so numerous and widespread in distribution that it is impossible to map them as separate features. Together with the intervening marshes and sandy lake flats, they constitute a complicated and undifferentiated relief.

The final chapter in the geologic history of the drainage basins has not yet been written. Numerous changes will, no doubt, be made throughout the years to come. The existing rivers will from time to time shift their courses and new streams will develop in the region. The drainage for the most part is still in a youthful condition but as the stage of maturity is approached and more tributaries are superposed upon the watershed slopes there is no question but that much of the peat and muck which now cover the surface will be removed. Extensive sandy flats will thus become exposed to renewed activity by the



Air photo showing oxbow lakes and canals in the lower reaches of the Manistiguo River.

wind and in time, the interstream areas will be covered with sand in place of the present muck and peat.

The low gradient in the lower reaches of the Manistique River has resulted in the development of broad meanders and cut-offs. (Plate XXVII) The sluggish character of this portion of the river was induced, in part at least, by damming of the older outlet by dunes formed along the Nipissing shore. Further changes in the outlet may have the effect of ponding tremendous volumes of water in the lower more or less unconfined valley ways.

In conclusion, it should be stated that the work which has so far been done in the area is inadequate to solve all of the problems which naturally present themselves in so complicated a region. The present paper is merely the first step in an attempt to unravel the events of the past. More exact data worked out on the basis of precise levelling are needed to correlate the various water planes accurately and to tie up definitely all of the steps involved in the lacustrine history. It is hoped, however, that the features which have been mapped will serve as a basis for more detailed studies of the local problems in the region.








FIGURE 16

LOCATION MAP
SHOWING
MAJOR FEATURES
OF THE
TAHQUAMENON AND MANISTIQUE DRAINAGE BASINS,
INCLUDING
ALGER, LUCE AND SCHOOLCRAFT COUNTIES.
S.G. BERGQUIST-GEOLOGIST.

BIBLIOGRAPHY

1. Hobbs, W. H., *Glacial Anticyclones*, The McMillan Company, 1926, pp. 198.
2. Gilbert, G. K., *Recent Earth Movements in the Great Lakes Region*, U. S. Geol. Survey, 18th Annual Report, Part 2, 1897. pp. 595-647.
3. Goldthwait, J. W., *The Abandoned Shore Lines of Eastern Wisconsin*. Wisc. Geol. and Nat. Hist. Survey, Bull. 17, 1907. 134 pp.
4. Fairchild, H. L., *Post-glacial Uplifts of Northeastern North America*, Bull. Geol. Soc. Am. Vol. 29, 1918. pp. 187-238.
5. Spencer, J. W., *The Focus of Post-glacial Uplifts North of the Great Lakes*, Jour. Geol. Vol. 19, 1911. pp. 57-60.
6. Taylor, F. B., *Deformation of Shore Lines*, U. S. Geol. Survey, Mon. 53, 1915. pp. 502-518.
7. Wisler, C. O., *Report on the Water Resources of the Tahquamenon River in Chippewa and Luce Counties*. Rept. of the Land Ec. Survey, May, 1927. 5 pp.
8. Scot, I. D., *Inland Lakes of Michigan*, Mich. Geol. and Biol. Surv., Pub. 30, Geol. Ser. 25, 1920. pp. 383.
9. Stevenson, Ellen B., *The Dunes of the Manistique Area*, Papers of the Michigan Academy of Science, Arts and Letters, Vol. 14, 1930, p. 481.
10. Bergquist, S. G., *The Occurrence of Glauconite in the Hermansville Formation of Alger County*, Papers of the Mich. Acad. of Science, Arts and Letters, 1929, pp. 231-237.
11. Davis, W. M., *Sub-glacial Origin of Eskers*, Proc. Bost. Soc. Nat. History, Vol. 25, 1892, pp. 477-499.
12. Trowbridge, A. C., *The Formation of Eskers*, Proc. Iowa Acad. Science, Vol. 21, 1914, pp. 211-218.
13. Russell, I. C., *The Malaspina Glacier*, Jour. Geol., Vol. 1, 1893, pp. 219-245.
14. Anderson, S. A., *The Waning of the Last Continental Glacier in Denmark as Illustrated by Varied Clays and Eskers*, Jour. Geol., Vol. 39, 1931, pp. 609-624.
15. Flint, R. R., *The Stagnation and Dissipation of the Last Ice Sheet*, Geogr. Review, Vol. 19, 1929, pp. 256-289.
16. Russell, I. C., *Drumlin Areas in Northern Michigan*, Bull. Geol. Soc. America, Vol. 16, 1906, pp. 577-578.
17. Chamberlain, T. C., *Proposed Genetic Classification of Pleistocene Glacial Formations*, Jour. Geol., Vol. 2, 1894, pp. 517-538.
18. Leverett, Frank, *Moraines and Shore Lines of the Lake Superior Region*, U. S. Geol. Surv., Prof. Paper 154A, 1928, p. 64.
19. Ibid.—p. 44.
20. Ibid.—p. 53.
21. Thwaites, F. T., *The Origin and Significance of Pitted Outwash*, Jour. Geol., Vol. 34, 1926, pp. 308-319.
22. Ibid.
23. Leverett, Frank, l. c., p. 53.
24. Leverett, Frank, *Surface Geology of Michigan* Geol. and Biol. Surv., Pub. 25, Geol. Ser. 21, 1917, p. 70.
25. Ibid.—pp. 66-67.
26. Winchell, N. H., *The Glacial Features of Green Bay of Lake Michigan, with some Observations of a Probable Former Outlet of Lake Superior*, Am. Jour. Sci., 3rd Ser., Vol. 2, 1871, pp. 15-19.
27. Bergquist, S. G., *The Occurrence of Glauconite in the Hermansville Formation in Alger County*, Papers of the Mich. Acad. of Science, Arts and Letters, 1929, pp. 231-237.
28. Leverett, Frank, *Moraines and Shore Lines of the Lake Superior Region*, U. S. Geol. Surv., Prof. Paper 154A, 1928, p. 67.
29. Ibid. p. 64.

30. Alden, W. C., Chicago Folio 81, Geol. Atlas U. S. Geol. Survey, 1902, p. 7.
31. Taylor, F. B., Leverett, Frank—The Pleistocene of Indiana and Michigan and the History of the Great Lakes, U. S. Geol. Surv. Mon. 53, 1915, p. 410.
32. Reudemann, R., and Ehlers, G. M., "Occurrence of the Collingwood Formation in Michigan." Contributions from the Museum of Geology, University of Michigan, Vol. 2, No. 2.
33. Leverett, Frank, U. S. Geol. Surv., Prof. Paper 154-A, 1928, p. 68.
34. Ibid—p. 70.
35. Hobbs, W. H., "The Late Glacial and Post Glacial Uplift of the Michigan Basin", Mich. Geol. and Biol. Surv. Pub. 5, Geol. Ser. 3, pp. 45-68.
36. Stevenson, Ellen B., The Dunes of the Manistique Area, Papers of the Michigan Academy of Science, Arts and Letters, Vol. 14, 1930, pp. 475-485.
37. Ibid—p. 483.
38. Bucher, W. H., Giant Ripples as Indicators of Paleogeography, Proc. Nat. Acad. Sci., Vol. III, 1917, pp. 285-291.
39. Udden, J. A., Mechanical Composition of Wind Deposits, Augustana Library Publication, No. 1, 1898.
40. Bryan, Kirk, Wind-Worn Stones or Ventrifacts—A Discussion and Bibliography. Reprint and Circular Series of the National Research Council, No. 8, 1928-1930, pp. 29-51.
41. Free, E. E., The Movement of Soil Material by the Wind, with a Bibliography of Eolian Geology by Stuntz, S. G., and Free, E. E., Bull. 68, U. S. Bur. of Soils, 1911, pp. 173.
42. Solger, F.,—Dünenbuch, Enke, Stuttgart 1910, pp. 404.
43. Högbom, Ivar,—Ancient Dunes of Northern and Middle Europe, Geogr. Annaler, V Bd, 1923, pp. 117-120.
44. Hobbs, W. H., The Glacial Anticyclones: the Poles of Atmospheric Circulation, The McMillan Company, 1926, 198 pp.
45. Antevs, Ernst, The Recession of the Last Ice Sheet in New England, Am. Geogr. Soc. Research, Ser. 11, 1922.
46. Reeds, C. A., Weather and Glaciation, Bull. Geol. Soc. Am., Vol. 40, 1929, pp. 597-630.

INDEX

	A	Page
Adams Trail, Munising		79
Aeolian, Activity (air currents and wind)	44, 53, 106, 113, 135	
Aerial photos, use of		47, 137
Aftonian interglacial stage, discussion of		20
Agricultural possibilities of the Manistique basin		65
of the Tahquamenon Valley		54, 58
Alden, W. C., reference to		93
Alger County, dunes in		113
glacial features in	61, 65, 71, 77, 78, 79, 81, 82, 86, 93, 99, 104, 113	
Lake Nipissing, shore lines in		104
Swamps in		43
Algoma stage of Great Lakes		110
Algonquin, Lake, discussion of	84, 44, 88, 95, 96, 99, 101, 112	
Alluvial fans, deposition of		82
Anderson, S. A., reference to		66
Anna River, outlet of Lake Algonquin		91, 133
Anteys, Ernst, reference to		133
Ape man of Java, time of appearance		20
Arkona Lake, discussion of		30
Artesian wells near Newberry		75
AuSable Lake, dunes west of		118
AuSable Point, Alger County, moraine near		78
AuTrain, dunes near	106, 121	
AuTrain Falls		91
AuTrain Lake, location of		106
AuTrain River, development of	91, 121	
dunes at mouth of		121
AuTrain-Whitefish Valley, outlet of Lake Algonquin		89
divide		91

B

Battle Creek and glacial drainage		26
Battlefield Beach of Lake Algonquin	93, 95, 97, 99	
Beaches of glacial lakes, records of		28
Beaver Lake, and Munising Moraine		78
dunes near		119
Belted lowlands, glaciation in	41, 93, 132	
topography of	43, 132	
Betsey lake, moraine near		82
reference to		57
Bibliography		139
Black River, a glacial drainage way		30
Black River limestone, in drift		66
Blaney, segment of Newberry Moraine, description of		73
Blind Sucker River, reference to	55, 99	
Blueberries in dry bogs		58
Bodi Lake, reference to		56
Bogs, quaking in Tahquamenon Basin		58
Boot Lake, location of		85
Boot Lake Plain, discussion of		85
Brule River, glacial lake outlet		88
Bryan Kirk, reference to		118
Bucher, W. H., reference to		118

C

Cambrian-Ozarkian contact locality	66, 80, 97	
Cambrian sandstones, at Upper Falls		51
cliffs of		104
in drift		66
near AuSable Point		119

	Page
Campsites, locations	54, 56
Case, Dr. E. C., whale skeleton discovered by	36
Chamberlain, T. C., reference to	73
Champlain marine stage, discussion of	36
Chapel Beach, dunes near	121
Charlotte moraine, position of	26
Chatham, drumlins near, description of	68, 80
Cheboygan moraine, position of	27
Chippewa County, Glacial features in	51, 56
Munising moraine in	80
Clear Lake, location of	85
Cold Spring, moraine near	80
Collingwood formation, locality of	96
"Complex," a land feature, defined	123
Continental uplift, effect of	36, 37, 44, 53, 54, 61, 109, 122, 134
Cook Mill, striae near	69
Cook's Moraine, described; location of	68, 101
Cordilleran glacier, defined	17
"Corrugated" relief, described, defined	125
Crisp Point, dunes near	122
moraine, described	81, 88
shore features on	97
Cromagnon civilization, time of	21
Cusino swamp, described	78

D

Danaher, divide near	92, 123
Davis, W. M. reference to	66
Dead Sucker River, valley of	55, 99
Defiance moraine, location of	27
Dollarville, glacial lake outlet near	92
moraine near	79
Dowagiac lake, formation of	26
Drainage channels of Great Lakes	26, 28, 29, 30, 31, 32, 33, 35, 36, 51, 63, 88, 89, 91, 92, 93, 95, 97, 99, 134
of Ice lobes, discussion of	25, 26
Drei-kanter, produced by sand blasts	118
Drift, character of in Trenary plains	65
glacial, definition of	20
Driggs creek, reference to	61
Drumlins, discussion of	68
near Eben Junction	65
of Menominee County	63, 68
of Trenary till plain	63, 68
Dune-like ridges, discussion of location	54, 97, 106
Dunes and wind work	106, 113, 115
cause of in Manistique valley	131
development of	44, 106
effect of on surface drainage	59, 122
in development of Tahquamenon drainage	54, 123, 131
in Manistique complex	123, 125
in Seney region	123, 125
inland, discussion of	129
Manistique River outlet blocked by	59, 131
near AuTrain	106
East Bay	119
Grand Marais	119
Grand Sable Lake	113
Lake Michigan	130
Lake Superior	54, 113
AuSable Lake	118
Beaver Lake	119

	Page
Dunes of Manistique Basin	59, 123, 131
swamp	129
valley	55, 59
Sucker basin	119
Tahquamenon basin	53, 97, 123, 131
the marshy lowlands	122
topography of, near Grand Marais	115

E

East Bay, dunes near	119
East Branch Fox River, discussion of	59, 61
Eben Junction, drumlins near	65, 68
Emmet Moraine, location of	27
Ehlers, G. M., reference to	96
Eskers, definition of	19
of Trenary till plain, discussion	63, 65, 66

F

Fairchild, H. L., reference to	44
Falls Au-Train near Forest Lake	91
of Tahquamenon	49, 51
Finger Lakes, N.Y. and Great Lakes	32
Fire Tower Hill, Seney, shore features near	95, 126
Flint, R. R., reference to	66
Flint moraine, location of	26
Flint river outlet of glacial lake	29
Flint-Mayville moraine and Imlay outlet	30
Fore dunes of Lake Algoma	110, 112
Forest cover type in Manistique Basin	125
near Seney	125
Forest, buried in sand dune area	115
Forests, pine, near Seney	125
Fort Brady Beach of Lake Algonquin	93, 95, 99
shore, description of	99
Fort Wayne outlet of Lake Maumee	28
Fort Wayne-Wabash moraine, position of	26
Fosse Lakes, description of	56
Fowler moraine, location of	26
Fox River, discussion of	59
Free, E. E., reference to	122

G

Gilbert, K. G., reference to	44
Gimlet Creek, dunes near	97
Glacial ancestors of Great Lakes	27-39
ice lobes, discussion of	22-23
Glacier, activity of in Tahquamenon-Manistique Basin	63, 132
developments of in North America	17
Gladwin moraine, location of	26
Gogebic Lake, outlet of a glacial lake	88
Goldthwaite, J. W., reference to	44
Goose Lake, a fosse lake	56
Grand Ledge moraine, location of	26
Grand Marais, beaches near, glacial features near	35, 95, 97, 99, 104
dunes near	113
Grand Portal headland, dunes near	121
Grand River, development of	27, 29, 30
Grand Sable bluffs, description of	119
Grand Sable Dunes, discussion of	113
Grand Sable Lake, glacial features near	113, 115
Great Lakes, history of	27
Green Bay ice tongue and drumlins near Chatham	80
Green Bay Lobe, glacial features developed by	63, 68, 69, 80

	Page
Green Bay Lobe, discussion of	63, 75, 80, 132
Greenland, glaciers of	17
Ground moraine, defined	19
Ground moraines and Munising moraine	77
Gulliver Lake, a remnant of glacial lake	108

H

Heidleberg man, time appearance of	20
Hermansville limestone, presence of	66, 91, 97
Hiawatha segment of Newberry moraine, discussion	61, 71
Hinge line and lake beach tilting	35, 36, 37
History, glacial, of Michigan	17
Hobbs, W. H., reference to	43, 47, 110
theory of wind direction from ice caps	130
Högboom, Ivar, classification of dunes by	129
Huron-Erie, ice lobe, discussion of	23

I

Ice accumulation, centers of	17
contact borders	77, 78
invasions, discussions of	18, 20
Illinoian drift described	20
ice invasion, discussion	20
Imlay outlet, discussion of	29
Indian Lake, discussion of and features near	59, 68, 108
Indian River, channel of	71
Indiana, moraines in	23, 24
Inland dunes, discussion of	129
Inland Steel Company Harbor, dune ridges near	125, 130
"Inner" morainic system, discussion of	75
Jonia moraine, location of	26

J

James Bay, ice center near	17
----------------------------------	----

K

Kalamazoo moraine, discussion of	25
Kankakee drainage	23
torrent, location of	25
Kansan glaciation discussed	20
Keewatin Glacier defined	17
Kingston Lake, plain bordering	86
Kingston Plain, description of	86
Kirkfield outlet of Great Lakes	35, 36, 44, 93, 97

L

Labradorean glaciation defined, work of	17
Lacustrine (lake) activity, discussion of	88, 133
Lag, gravel, defined	118
Lake Algoma, discussion of	110
Lake Algonquin, discussion of	34, 44, 88, 89, 95, 99, 101, 126, 133
Lake Arkona, discussion of	30
Lake AuTrain, description of	106
Lake Border morainic system, location of	26, 27
Lake Chicago, development of	29, 31, 34
Lake Dowagiac, discussion of	26
Lake Duluth, discussion of	31, 34, 88
Lake Huron, beginning of	30
Lake Huron ice lobe	23
Lake Lundy, discussion of	33
Lake Maumee, discussion of	28
Lake Michigan, beginning of	31

	Page
Lake Michigan, dunes of, ice lobe	130
Lake Michigan, ice lobe	23, 26, 132
Lake Nipissing, development of	104
Lake Ontonagon, discussion of	88
Lake Saginaw, discussion of	30
Lake Superior, ice lobe, moraines of	23, 71, 75, 76, 80, 81
dunes of	113
Lake Warren, discussion of	33
Lake Wayne, discussion of	32
Lake Whittlesey, discussion of	30
Lakes, fosse, description of	56
morainal, description of	56
of the Manistique drainage system	59
of the Tahquamenon drainage basin	56
pit, description of	56
recreational value of in Luce County	56
shores of former lakes—See glacial lake discussions.	
Land Economic Survey referred to	47
Lansing moraine, location of	26
Leverett, Frank, quoted	77, 87, 88, 91, 104
reference to	21, 28, 47, 65, 77, 78, 85, 87, 88, 91, 92, 99
Leverett, F., and Taylor F., reference to	93
Lilley Lake, location of	85
Little Two Hearted River, description of	55
Lookingglass River, development of	27
Lower Falls of Tahquamenon	51
Lower Algonquin shores in Tahquamenon region	96
Lowest Algonquin shores	99
Luce County, dunes in	121
glacial features in	49, 53, 54, 55, 57, 73, 74, 79, 81, 82, 86, 95, 96, 97, 99, 104, 121
lakes of, recreational value of	57, 134
Niagara escarpment in	51
Lyons moraine, location of	26

M

Manistique, location of on glacial shore	59, 108
shore features near	106, 108, 123, 130
Manistique Basin, discussion of features of	43, 59, 101, 123
Manistique Complex described	123, 125
Manistique Lake, discussion of	56, 59, 74
Manistique River, discussion of	43, 59, 101, 108, 131, 132, 134
Manistique swamp	59, 92, 125, 129
Maple River, a glacial outlet	27, 30
Marseilles moraine and Kankakee Torrent	25
Marsh Creek, reference to	61
Marshes of Luce County	57
Maxinkuckee moraine, location of	25
McDonald Lake, glacial remnant	108
McMillan, location of	53, 87
McMillan Straits, and establishment of present drainage	51, 93
glacial shore features near	92, 95
Melstrand, character of moraine near	85
Michigan ice lobe, discussion of	63, 69, 132
Milakokia River, dune ridges near	106, 125
Mississinewa Moraine, position of	25
Mohawk Outlet of Great Lakes	32, 34
Moraines, control of rivers by	133
definition of	19
ground (till plains) defined	19
in Michigan, development of	23
of Michigan and Superior lobes	26, 71, 75, 77
of Munising system	75, 77
Mount Clemens, Moraine, location of	27

	Page
Muck lands, character of	53, 54, 135
Muck in Manistique swamp	59, 129, 131
in Luce County	54
Mud Lake, reference to	57
Munising Inner Morainic system	75, 97
Munising, Algonquin shoreline near elevation	35
location of on Nipissing terrane	106
moraine near	77
Munising moraine, discussion of	75, 80, 83, 91, 92, 97, 132
in Alger county	77, 79, 80, 99
in Luce County	79, 80
in Chippewa County	54, 80
in Schoolcraft County	79, 80
outwash plains associated with	83
significance of, discussion of	80
Murray Lake, a fosse lake	56
Muscalonge Lake, a pit lake	56
Muskeg, marsh, and swamps, discussion of	57, 58

N

Nadoway Point, terminus of Newberry moraine	80
Nebraskan glaciation, beginning of	20
Newberry, deep well at asylum	75
glacial shore features near	95
location of	53
morainic area, south of	53
municipal water supply of	75
Newberry moraine, discussion of	61, 69, 81, 132
control of rivers by	133
Newberry-Munising moraine, merger of	80
Niagara escarpment	51, 73, 74
and Blaney moraine	74
Niagara Falls, development of	34, 36
Nipissing beaches, tilting of	37
Nipissing Great Lakes, discussion of	36, 104
Nipissing shore features in Lake Michigan region	106
Superior region	73, 104
Alger County	104
Luce County	104
North Bay outlet of glacial lakes	92, 95, 104
North Manistique Lake discussion	56, 74

O

Ohio River relation of to ice border	17, 20
Old shores in Grand Marais region	99
Ontonagon, glacial lake	88
Ottawa drainage channel, outlet of Great Lakes	88
"Outer" Morainic system, discussion of	69
Outwash plains, discussion of	19, 82
of Cook's moraine	69
Owosso Moraine, location of	26

P

Peat bogs, discussion of	57, 129, 130, 131
Peat, formations of	129
marshes between dunes	126
Pike lake, reference to, a pit lake	56
Pithecanthropus erectus, time of appearance	20
Pit lakes, description of	56
Pleistocene Epoch defined	17
Poindexter, F. O., reference to	108
Point Aux Barques, dune barrens near	130

TAHQUAMENON-MANISTIQUE DRAINAGE BASIN

147

	Page
Port Huron moraine, discussion of	27
Portland moraine, location of	26
Post-Nipissing shore features, discussion of	108

R

Raisin River, Illinoian drift in bed of	20
Reeds, C. A., reference to	133
Resort sites, discussion of	56
on "ridges" of Luce County	54
Russell, I. C., reference to	66, 68

S

Sable Creek, channel, shifting of	115
Saginaw Bay, relation of glacier to	23, 26
Saginaw Lake, formation of	23, 30
Saginaw glacial lobe, discussion of	23
St. Johns moraine, discussion of	26
St. Joseph, River valley of	25
St. Lawrence River, development of as outlet	36
Sand dunes along Lake Superior	54, 113
in Tahquamenon valley	54, 97, 123, 131
Sand Lake, location of	85
Sangamon interglacial stage, discussion of	21
Scaffold Hill, storm beaches near	77
Schoolcraft County, glacial features in	68, 69, 78, 80, 83, 86, 87, 101, 108, 122, 125
Scott, I. D., reference to	47, 112, 122, 126, 130
Scott, I. D., and Stevenson, Ellen, studies of	112
Seney, dunes near	126
Seney Complex, discussion of	123, 125
Seul Choix Point headland, glacial features near	130
Sheldrake drainage system, discussion	55
Shore features of Lake Algonquin	95, 99
of Lake Nipissing	104
of post-Nipissing time	108
Smith, Dr. R. A., reference to	48
Soils for clover and alfalfa, description of	54
Solger, F., classification of dunes by	129
Spencer, J. W., reference to	44
Stanley, Geo., studies of	93
Steuben segment of Newberry moraine, description	71
Stevenson, Ellen, reference to and quoted	59, 110, 112, 130
Straits of Mackinac, opening of	35
Striae, glacial near Cook's Mill	69, 132
Sturgeon River moraine, Delta County, reference to	75
Sturgis moraine, location of	25
Sucker River and basin, glacial features of, dunes in	55, 56, 104, 120
Superior lobe, features developed by	43, 69, 80, 120, 133
Swamps, character, development, discussion, of	43, 44, 57

T

Tahquamenon drainage system, discussion of	43, 49, 51, 80, 92, 96, 133, 134
dunes of	54, 123, 131
glacial features of	43, 51, 96
morainic control of	133
Tahquamenon falls, description of	50, 51
Tahquamenon-Manistique outlet of Superior Basin	92
Tahquamenon swamp discussion	97
Taylor, Frank, reference to	44, 93
quoted	93, 99
Tekonsha moraine, location of	25
Temple Lake, referred to, dunes near	56, 106
The Great Lakes, history of	27

	Page
Thompson, lake shores near	108, 110
"Thumb" area, glacial lakes of	29, 30
Thwaites, reference to, quoted	86, 87
Till Plains, defined	19
Timber of Luce County, Seney and Manistique Complexes	58, 123
Trenary plains, discussion of	63, 66, 68, 132
Trent outlet of Great Lakes, discussion of	35, 44, 93
Trenton limestone, drift locality	66
in AuTrain-Whitefish valley	91
Trowbridge, A. C., reference to	66
Two Hearted Lakes, reference to	53
Two Hearted River, dunes near	54
Two Hearted River system, discussion of	54

U

Ubley, outlet, development of	30, 31
Udden, J. A., reference to	118
Uplifts, discussion of effects of	36, 37, 44, 53, 61, 109, 122, 134
Upper Falls of Tahquamenon	49, 51, 53, 134

V

Valparaiso Moraine, position of	26
Varves in Anna River Valley	133
Ventrifacts, defined	118

W

Wabash River Outlet of Lake Maumee	28
West Branch Moraine, location of	26
West branch of Fox River, reference to	61
of Manistique River	61
Wetmore Plain, discussion of	83
stone beaches near	77, 92
Whales in Michigan drift, fererence to	36
Whitefish Lake, discussion of	59
Whitefish River, development of	91
Winchell, N. H., reference to	89
Windwork, discussion of	44, 53, 106, 135
Wisler, C. O., reference to	51
Wisconsin glaciation, discussion of	20, 21
Wisconsin till, described	21

Y

Yarmouth Interglacial Stage, discussion of	20
Ypsilanti, Maumee Beach near	29

